



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

507 .U58 NO.55

C.1

A contribution to the

Stanford University Libraries



3 6105 046 413 287



201

458

SMITHSONIAN INSTITUTION
UNITED STATES NATIONAL MUSEUM

BULLETIN

OF THE

UNITED STATES NATIONAL MUSEUM

No. 55



WASHINGTON
GOVERNMENT PRINTING OFFICE
1905

ADVERTISEMENT.

This work (Bulletin No. 55) is one of a series of papers intended to illustrate the collections belonging to or placed under the charge of the Smithsonian Institution and deposited in the United States National Museum.

The publications of the National Museum consist of two series—the *Bulletin* and the *Proceedings*.

The *Bulletin*, publication of which was commenced in 1875, is a series of elaborate papers issued separately and based for the most part upon collections in the National Museum. They are monographic in scope and are devoted principally to the discussion of large zoological groups, bibliographies of eminent naturalists, reports of expeditions, etc. The bulletins, issued only as volumes with one exception, are of octavo size, although a quarto form, known as the Special Bulletin, has been adopted in a few instances in which a larger page was deemed indispensable.

The *Proceedings* (octavo), the first volume of which was issued in 1878, are intended primarily as a medium of publication for newly acquired facts in biology, anthropology, and geology, descriptions of new forms of animals and plants, discussions of nomenclature, etc. A volume of about 1,000 pages is issued annually for distribution to libraries, while a limited edition of each paper in the volume is printed and distributed in pamphlet form in advance.

In addition, there are printed each year in the second volume of the Smithsonian Report (known as the Report of the U. S. National Museum), papers, chiefly of an ethnological character, describing collections in the National Museum.

Papers intended for publication by the National Museum are usually referred to an advisory committee, composed as follows: Frederick W. True (chairman), George P. Merrill, Otis T. Mason, James E. Benedict, Walter Hough, T. W. Stanton, Leonhard Stejneger, and Marcus Benjamin (editor).

S. P. LANGLEY,

Secretary of the Smithsonian Institution.

WASHINGTON, U. S. A., December 1, 1905.

A CONTRIBUTION TO THE OCEANOGRAPHY OF THE PACIFIC

COMPILED FROM DATA COLLECTED BY THE UNITED
STATES STEAMER *NERO* WHILE ENGAGED IN
THE SURVEY OF A ROUTE FOR A
TRANS-PACIFIC CABLE


BY

JAMES M. FLINT

Medical Director, U. S. Navy; Curator, Division of Medicine, U. S. National Museum



Washington
Government Printing Office
1905



2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2

TABLE OF CONTENTS.

	Page.
Introduction	1
Plan of survey	2
Track	3
Depths	3
Gradients	5
Temperatures	5
Air and surface water	5
Bottom temperatures	7
Character of bottom	9
Deposits	10
Red clay	10
Globigerina ooze	11
Diatom ooze	12
Radiolarian ooze	13
Volcanic mud	13
Blue mud	13
Green mud	14
Record of the detailed examination of selected specimens of deposits from stations on the outward voyage of the Nero	15
(A) Honolulu to Midway Islands	15
(B) Midway Islands to Guam	17
(C) Guam to Luzon	21
(D) Guam to Yokohama, Japan	24
Abstract of the official record of soundings	27
Hawaiian Islands to Midway Islands	27
Midway Islands to Guam	30
Guam to Luzon	37
Luzon to Guam	40
Guam to Yokohama	42
Yokohama to Guam	46
Guam to Midway Islands	50
Midway Islands to Hawaiian Islands	57
Explanation of plates	61

82945

A CONTRIBUTION TO THE OCEANOGRAPHY OF THE PACIFIC

By JAMES M. FLINT,

Medical Director, U. S. Navy; Curator, Division of Medicine.

INTRODUCTION.

In the early part of the year 1899 the U. S. S. *Nero*, a steam collier of 4,925 tons displacement, which had been purchased for use during the Spanish-American war, was fitted out by the Navy Department, equipped with the necessary apparatus, and dispatched from San Francisco under the command of Commander Charles Belknap, U. S. Navy, with instructions to survey a route for a telegraph cable between the United States, the Philippine Islands, and Japan.

On account of the illness of Commander Belknap, he was relieved from command on the arrival of the ship at Manila by Lieut. Commander H. M. Hodges, U. S. Navy, who remained in charge of the survey until its completion.

Several previous surveys having established a satisfactory route between the coast of California and the Sandwich Islands, the actual work of the *Nero* began at Honolulu, from which port the ship sailed on the 6th day of May, 1899.

The following table of dates, distances, and number of soundings furnishes an abstract of the cruise:

Locality.	Date.	Distance run.	Number of soundings.
	1899.	<i>Knots.</i>	
Left San Francisco	Apr. 22		
Arrived Honolulu	May 2		None.
Left Honolulu	May 6		
Arrived Midway Island	May 22	1, 184. 5	195
Left Midway Island	May 24		
Arrived Guam	July 5	3, 520. 35	467
Left Guam	July 7		
Arrived Dingala Bay, Luzon	Aug. 1	1, 400. 15	191

Locality.	Date.	Distance run.	Number of soundings.
	1899.	<i>Knots.</i>	
Left Dingala Bay	Aug. 1		
Arrived Manila	Aug. 4		None.
Left Manila	Aug. 15		
Arrived Dingala Bay	Aug. 18		None.
Left Dingala Bay	do		
Arrived Guam	Sept. 7	2, 753	136
Left Guam	Sept. 9		
Arrived Yokohama	Sept. 24	1, 427. 70	248
Left Yokohama	Oct. 10		
Arrived Guam	Nov. 2	2, 941. 50	231
Left Guam	Nov. 12		
	1900.		
Arrived Midway Island	Jan. 3	5, 725. 30	402
Left Midway Island	do		
Arrived Honolulu	Jan. 29	2, 567. 15	204
Left Honolulu	do		
Arrived San Francisco	Feb. 11		None.
Total		21, 519. 65	2, 074

From the above table it appears that the distance sailed while on actual survey work was 21,519 geographical miles and the number of soundings 2,074, or an average of one sounding at every ten miles of distance run. Measuring the direct course of the survey, 6,144 knots, there are records of soundings averaging one every three miles of the route.

PLAN OF SURVEY.

The instructions regarding the survey were, in brief, to follow as nearly direct lines as practicable from Honolulu to Midway Island, thence to Guam, thence to Luzon, and also from Guam to Japan. Soundings were to be taken on the outward voyage at intervals of 10 and 2 miles alternately; temperatures of the air, surface and bottom of the sea to be recorded; currents noted; samples of bottom material brought up in a sounding cup to be preserved, etc. The return course was planned to cross the primary route zigzag at angles of 45°, the sides of the angle to be 20 miles in length; soundings to be taken at the apices of the angles. This plan was effectively carried out, modified somewhat in detail by circumstances, especially as regards intervals between soundings and detours from the main line in order to develop marked irregularities of the contour of the ocean bed. In this manner an examination was made of a belt of ocean about 14 miles wide and over 6,000 miles in length, unequalled in thoroughness, so far at least as soundings are concerned, by any survey hitherto made of an ocean tract.

TRACK.

It would be unnecessary for the purpose of this study to attempt to present in detail the exact courses followed by the vessel, which were at times quite erratic, in the effort to find the most feasible location for the cable. It is to be understood, therefore, that the accompanying track charts do not represent the exact courses of the ship nor the line determined upon for the cable or followed in the laying of it. The stations charted, however, are supposed to be accurately located. Before preparing the charts certain stations on the outward voyage were selected for careful examination of bottom material. The considerations governing the selection were chiefly the depth, the macroscopic appearances of the bottom specimens, and the distances. These stations were afterwards plotted and connected by a continuous line. In a few instances outlying stations of especial interest have been indicated on the charts, and a serial number and depth given for each. On the first or diagrammatic chart the station numbers included within each five degrees of longitude or latitude are given for both the outward and homeward voyages. By means of this index it is easy to locate approximately any station mentioned in the record.

DEPTHS.

The graphic representation of the contour of the ocean bed along the course of this survey is less simple and satisfactory than usual, because of the breadth of track explored and the large number of soundings recorded. It should be noted, in examining the contour charts appended, that the lines are drawn from the localities indicated on the track charts as stations selected for special examination of bottom material, and one of the principal determining factors in the selection was that of depth. Therefore, the contour charts may be said to represent the extremes of elevation and depression along the main line of the outward voyage only, without taking into account intermediate or outlying irregularities of surface. The omitted stations can be easily supplied from the record if greater detail is desired.

Leaving the island of Oahu of the Hawaiian group, the depth increases quite rapidly until it reaches about 2,500 fathoms. This depth is reached less than 30 miles due north of the island. From this point nearly to Midway Island there is a comparatively level plain, broken only by two or three outlying peaks from the mountain range whose highest summits show themselves as small islands or reefs a little to the westward of the line of survey. One of these peaks appears at station 93, where there is a sharp rise to 1,463 fathoms, which, however, as rapidly falls away to the normal level a few miles to the northward. A smaller projecting spur is indicated at station 106, 2,002 fathoms. Another sudden rise to 1,726 fathoms

appears at station 124, followed by depressions to more than 2,600 fathoms a few miles away, both to the westward and northward. With these exceptions the range of variation is practically between 2,500 and 3,000 fathoms for the whole distance until the immediate vicinity of the Midway Islands is reached.

About 30 miles to the southward and westward of Midway Island a very bold peak was discovered rising abruptly from the ocean floor, 2,000 fathoms below the sea level, to a height only 82 fathoms below the surface.

Passing from the vicinity of Midway Islands a nearly level plain is found, extending about 1,000 miles to the westward, upon which the extremes of depth of water are 2,926 and 3,382 fathoms. About mid-distance between Guam and Midway Islands what is apparently a mountain range is encountered, extending over 3° of longitude, with varying depths from 3,000 to 720 fathoms. On the westward side of this mountain range another plain below the 3,000-fathom line extends a distance of about 300 miles. From the western limit of this plain until Guam is reached the contour is quite irregular. Extensive detours both north and south of the direct course developed a mountainous region, with peaks rising to 689 fathoms below the sea level, and valleys descending to a depth of more than 5,000 fathoms. Four soundings below the 5,000-fathom line were made, with the record of 5,070, 5,101, 5,160, and 5,269 fathoms. These were in the abyss now known as the "Nero Deep." The last-named sounding was numbered 1488, in latitude $12^{\circ} 43' 15''$ north, longitude $145^{\circ} 49'$ east, about 75 miles east-southeast from the island of Guam, and is the deepest sounding ever recorded, being only 66 feet less than 6 statute miles.

From Guam to Luzon the ocean bed is for the most part gently undulating, at depths varying from 2,500 to 3,000 fathoms. About 120 miles west of Guam (station 688) there appears a sharp elevation to 1,346 fathoms, which however soon subsides to the normal depth of about 2,700 fathoms. Again, about 600 miles from Guam (station 760) a rise to 1,560 fathoms is encountered. From the data at hand this latter would seem to be a peak rather than a mountain range, since soundings east, west, and south show speedy subsidence to nearly normal depths. At station 784 the depth reaches 3,547 fathoms, with several soundings in that vicinity below 3,000. Approaching Dingala Bay on the east coast of Luzon and about 120 miles distant (station 864) another peak appears with summit only 821 fathoms below sea level. In this instance also, soundings north, east, and west soon develop normal depths.

From Guam to Yokohama the route lies to the westward of the Ladrone Islands and to the eastward of the Bonin group. For a distance of 500 miles or more from Guam the soundings show a gently undulating plain at an average depth of about 2,100 fathoms. Between

latitude $21^{\circ} 45'$ and $22^{\circ} 8'$ north and longitude $143^{\circ} 45'$ and $143^{\circ} 20'$ east three sharp peaks arise along a line about 35 miles in length and northwesterly in direction. On the first, or most southerly, the sounding record is 483 fathoms; on the second, about 18 miles away, the record is 838 fathoms; and on the third, 20 miles farther to the northwest, 802 fathoms. There are valleys 1,000 fathoms deep between these peaks. The indications point to a continuous range of mountains connecting the Ladrone Islands with the Bonin group. After dropping down the eastern slope of the above-mentioned peaks, the depth increases by an easy gradient to 3,595 fathoms at station 1095, rising and falling gently until at station 1126 a sounding of 972 fathoms locates an outlying spur from the Bonin range. Still farther to the northward and westward, at station 1135, the bottom drops to 3,421 fathoms, followed by gentle slopes up to 1,500 and down to 2,900 fathoms, until the Gulf of Tokyo is reached.

GRADIENTS.

In computing the gradients from station to station serially on the outward voyage only, involving 1,100 soundings, sixty-nine localities only are found where the gradient exceeds 10 per cent. These higher grades are for short distances only, averaging less than 5 miles, and confined to a few regions, especially to the vicinity of Midway Islands, Guam, and the mountain range halfway between the above-mentioned islands. Of the sixty-nine localities showing a grade above 10 per cent, fifty have an incline between 10 and 20 per cent, eleven between 20 and 30 per cent, and six between 30 and 40 per cent. At the entrance to Port Tarafoto, on the east coast of Guam, two soundings one-fourth of a mile apart show a difference of depth of 123 fathoms, equivalent to a gradient of about 51 per cent. Also on the declivity of the peak southwest of Midway Islands, which rises to 82 fathoms beneath the surface of the water, there is a change of depth of 1,269 fathoms (7,614 feet) in a horizontal distance of 1.8 sea miles, a gradient of 70 per cent. With these few and localized exceptions the bed of the Pacific Ocean, as developed by this survey, though rising here and there near to the sea level, and again descending to depths of 5 or 6 statute miles, follows easy gradients. On the great plain to the westward of the Midway Islands, 1,000 miles in breadth, the average gradient is less than 1 per cent—in one instance only rising to 4.5 per cent, for a distance of 2 miles.

TEMPERATURES.

AIR AND SURFACE WATER.

The temperature of the air on board the ship, and of the water near the surface, was taken at nearly all the sounding stations. These stations numbered, on the average, about ten each day on the outward

voyage, and eight on the return voyage, distributed at nearly equal intervals over the twenty-four hours.

The following table presents the results of certain computations from the official records. (All temperatures are given in degrees Fahrenheit.)

Temperature of air and surface water.

Locality.	Date.	Number of observations.	Air.			Water.		
			High.	Low.	Average.	High.	Low.	Average.
Hawaiian Islands to Midway.	May 6 to May 24, 1899.	187	81	66	73.3	78	67	73.2
Midway to Hawaiian Islands.	Jan. 3 to Jan. 29, 1900.	185	79	61	69.1	77	65	72
Midway to Guam ...	May 24 to July 6, 1899.	463	92	72	79.2	86	70	80.6
Guam to Midway ...	Nov. 12, 1899 to Jan. 1900.	405	87	63	77.5	85	66	80.6
Guam to Luzon	July 7 to Aug. 1, 1899.	191	90	75	82.6	89	82	84.3
Luzon to Guam	Aug. 19 to Sept. 9, 1899.	134	91	77	82.8	87	80	84.5
Guam to Yokohama.	Sept. 9 to Sept. 24, 1899	248	90	68	81.2	87	75	83.8
Yokohama to Guam.	Oct. 10 to Nov. 2, 1899.	228	91	67	79	86	70	81.9

It will be seen from the above table (1) that the average temperature of the air, in these regions uninfluenced by the proximity of other than small and scattered islands, varies little from that of the contiguous waters of the sea. (2) That in the region between the Hawaiian Islands and Guam the difference between summer and winter temperatures of both air and water is quite small. On the round trips between Guam and Luzon and Guam and Yokohama, each having occupied only about two months, there are not sufficient data for estimation of seasonal changes in these regions. In considering extremes of temperature, it should be remembered that Midway Islands and Yokohama are both in considerably higher latitudes than the Hawaiian Islands, Guam, and Luzon, and, other conditions being equal, the lowest temperatures would naturally be found in the higher latitudes.

Thus, the surface temperature in the vicinity of Oahu is about $75^{\circ}.4$, while in the vicinity of Midway it falls to about 71° . Leaving Midway with an average of $70^{\circ}.4$ at the first eleven stations, the surface temperature rises to an average of $84^{\circ}.8$ at the last thirty-one stations approaching Guam. From Guam until within 300 miles of Yokohama the surface water remains near 85° , falling to an average of $76^{\circ}.7$ at the last eighteen stations.

The diurnal variations of temperature were of course greater in the air than in the water near the surface. The normal range of variation was from 4° to 7° for the air, and 1° to 3° for the surface water. The

extreme range of air temperature for any one day was 14° , January 17, 1900. There is also one record of 13° August 30, 1899, four of 11° , and four of 10° . Averages are shown in the following table:

Average daily variations.

Locality.	Date.	Air.	Surface water.
Hawaiian Islands to Midway	May 6 to May 22	5.2	2.4
Midway to Hawaiian Islands	Jan. 3 to Jan. 29	4.7	1.9
Midway to Guam	May 24 to July 5	5.8	2.0
Guam to Midway	Nov. 12 to Jan. 3	3.4	1.1
Guam to Luzon	July 7 to Aug. 1	5.7	1.7
Luzon to Guam	Aug. 18 to Sept. 7	7.7	2.6
Guam to Yokohama	Sept. 9 to Sept. 24	5.3	2.1
Yokohama to Guam	Oct. 10 to Nov. 2	5.8	2.2

How much the recorded air temperatures may have been affected by local conditions, such as radiation from the heated deck at midday, or evaporation from a wet deck, it is impossible to estimate.

BOTTOM TEMPERATURES.

No serial temperatures were taken. Observations of bottom temperatures on both outward and homeward voyages to the number of 604 are reported. In drawing conclusions from the records of these observations, some allowance should be made for the difficulties attending the measurement of temperatures at great depth, because of the delicacy of the instruments, the enormous pressures to which they are subjected, the shocks to which they are liable, and the vibration tending to displace the index as the thermometer is drawn up. Professor Tate says: "The circumstances under which thermometers are let down and drawn up again at sea are extremely unfavorable to accuracy of observation." In the column of remarks, on the *Nero* records, it is repeatedly noted that "Thermometer failed to work." So that where striking variations from normal temperatures, at given depths and in neighboring localities, appear on the record, the probabilities seem largely in favor of the assumption of instrumental, or possibly clerical, errors rather than of great eccentricities of temperature, unless there should appear to be something in the local conditions reasonably to account for the variation.

^a Results of the Exploring Voyage of H. M. S. *Challenger*.

The following table presents an abstract of the records of bottom temperatures:

Depths.	Number of observations.	High.	Low.	Average.
		°	°	°
Less than 500 fathoms	1			43.7
500 to 600 fathoms	1			39.8
600 to 700 fathoms	3	40.5	38.3	39.4
700 to 800 fathoms	5	38.6	36	37.3
800 to 900 fathoms	7	41.1	36.7	38
900 to 1,000 fathoms	3	37	36	36.4
1,000 to 1,500 fathoms	42	38	35	35.46
1,500 to 2,000 fathoms	83	39	35	35.31
2,000 to 3,000 fathoms	^a 266	36	34.2	35.17
3,000 to 4,000 fathoms	^b 188	36.3	34	35.22
4,000 to 5,000 fathoms	3	35.6	35.4	35.50
5,070 and 5,101 fathoms	2	36	35.9	35.95

^a 16 records thrown out.

^b 10 records thrown out.

The high temperature average, between 800 and 900 fathoms, is due to the exceptional record of 41°.1 at station 1225, in immediate proximity to the volcanic island of Oshima or Vries Island, at the entrance to the Gulf of Tokyo; also two records of 39° at stations 1569 and 1570, on the summit of a high peak or ridge about 450 miles to the eastward of the island of Guam. The average of the other four records is 36°.97.

In the series of observations at depths between 1,000 and 1,500 fathoms the is record of 38° at station 1678, and 37°.3 at the adjoining station 1677. These two stations are on one of the peaks of the mountain range in midocean between Midway and Guam. There are no other records of temperature above 37° at these depths.

Only two stations between the 1,500 and 2,000 fathom line record temperature above 37°, namely: Stations 1000, 39°.3 and 1009, 37°.3, about 60 and 120 miles, respectively, to the northward of Guam. There is probability of error in one or both of these observations.

In making up the average of temperatures between 2,000 and 3,000 fathoms, 16 of the 266 observations have been omitted from the calculations. In some of these cases "incorrect" is noted on the original record; in others, the probability of instrumental or clerical error is so much greater than the probability of existence of local conditions capable of producing such deviations from the normal range of temperature as to justify their exclusion. The omissions are stations 131 (44°), 138 (51°.7), 140 (44°.8), 232 (39°.4), 243 (38°.2), 477 (38°.8), 479 (44°.5), 719 (67°.6), 722 (67°), 723 (67°), 962 (37°), 1508 (33°), 1511 (33°), 1512 (34°), 1513 (34°), 1514 (34°). The last five of these rejected observations were taken by a thermometer concerning which it is noted: "Correction not known." This thermometer

being replaced by another, the temperatures are again recorded at the normal of 35° and above.

Of the 188 temperatures taken at depths from 3,000 to 4,000 fathoms, 10 have been excluded from the computation of averages, for the reasons given above. They are the following: Stations 239 (52°), 244 (38°), 251 ($38^{\circ}.9$), 257 (60°), 312 ($38^{\circ}.4$), 422 ($37^{\circ}.8$), 489 ($40^{\circ}.2$), 501 ($37^{\circ}.4$), 790 (32°), 809 ($22^{\circ}.9$).

Three temperature observations were made between 4,000 and 5,000 fathoms, and two at depths of 5,070 and 5,101 fathoms respectively, all in the abyss southward and eastward of Guam.

The obvious inference from the above computation is that the temperature of that part of the Pacific Ocean covered by this survey falls rather rapidly from the surface to about 600 fathoms, then very slowly to about 2,500 fathoms, where the normal temperature varies but slightly from 35° F. Below 2,500 fathoms there appears to be a slight rise of a fraction of a degree. But it is open to question if this apparent rise may not be due to the effect of the enormous pressure of three to five tons to the square inch, at these great depths, upon the instruments.

CHARACTER OF BOTTOM.

The character of the bottom indicated on the record by abbreviations, refers only to the gross appearances of the material recovered in the sounding cup, when fresh from the water. Translated into the terms of the usual scientific classification, the brown mud ("br. m.") of the record is generally the red clay of the oceanographer, or rarely volcanic mud from deep water. With few exceptions what is designated coral sand ("co. s.") is globigerina ooze. The rock (R. or r.) has, in every case examined, proved to be fragments of pumice or manganese-iron concretions; the black specks also are almost always particles of manganese iron. Except in the immediate vicinity of a shore, gravel (G. or gvl.) is, in this part of the ocean at least, coarse volcanic débris which has been distributed by wind or wave all over the sea, and has finally found its way to the bottom. The sand (S. or s.) so often noted consists of finer mineral particles from the same source as above-mentioned, except near the shores of islands.

Other abbreviations than those just given, used in the columns for character of bottom, refer to color, size, etc.—bk=black; br=brown; dk=dark; gy=gray; lt=light; rd=red; wh=white; y. or yl=yellow; crs=coarse; fn=fine; hrd=hard; rky=rocky.

DEPOSITS.

The accepted classification of marine deposits, by Dr. John Murray and Dr. A. F. Renard,^a is as follows:

Marine deposits.

1. Deep-sea deposits beyond 100 fathoms.	Red clay.	} Pelagic deposits formed in deep water removed from land.
	Radiolarian ooze.	
	Diatom ooze.	
	Globigerina ooze.	
	Pterapod ooze.	
	Blue mud.	} Terrigenous deposits formed in deep and shallow water close to land masses.
	Red mud.	
	Green mud.	
	Volcanic mud.	
	Coral mud.	
2. Shallow-water deposits between low-water mark and 100 fathoms.	Sands, gravels, muds, etc.	
3. Littoral deposits between high and low water marks.	Sands, gravels, muds, etc.	

Only 22 soundings are recorded within the 100-fathom line, and from several of these no specimens have come to hand. Practically, therefore, only deep-sea deposits have to be considered in this report.

Red clay.—Of the above-mentioned classes of deposits by far the most extensive is red clay. This, as it appears in the specimens received, is a smooth, sticky mud, varying in color from light yellowish-brown (fawn color) to dark chocolate, these colors being somewhat modified in individual instances by exposure to light, and especially by drying. In composition it consists of (1) extremely fine, amorphous particles of clayey matter, mostly hydrated aluminum silicate and the débris of other minerals; (2) the remains of calcareous organisms (foraminifera, coccospheres, and rhabdospheres), this constituent, however, rapidly disappearing at depths of about 2,500 fathoms; (3) siliceous organic remains (sponge spicules, radiolarian skeletons, and the frustules of diatoms; (4) mineral fragments, mostly of volcanic origin, at least in this part of the ocean; and (5) certain products of local chemical reactions, especially nodules, coatings, and grains of manganese peroxide, crystals of phillipsite, and particles of palagonite. The proportions of these constituents vary greatly along the line and even from station to station. As has been stated, foraminifera disappear, for the most part, at depths below 2,500 fathoms; radiolaria are likely to be more numerous in the deeper waters; diatoms are nearly everywhere, but only occasionally in great numbers. Mineral fragments may be so minute in some specimens that they pass over almost

^a Report on Deep Sea Deposits, based on specimens collected during the voyage of *U. S. Challenger*.

entirely in the fine washings, while in others they may be comparatively coarse. Volcanic glass is sometimes present in notable quantity. Manganese-iron nodules, and concretions upon other minerals, are almost universally present. They are the black specks ("bk. sp.") so frequently recorded on the official records, the larger ones being generally referred to as rock ("R.") Phillipsite is a frequent constituent. It is found as quite perfect crystals, single, twinned, or multiple, or more frequently as spherules made up of crystals arranged radially. The simpler forms are found in great numbers at station 331 (2,997 fathoms), and the spherules at station 495 (3,204 fathoms). Vertebrate remains, teeth of sharks and other fishes, and otoliths, have not been observed in this or other deposits, though carefully looked for.

This red clay deposit is indicated in 75 per cent of the soundings from which specimens were received (1,043 out of 1,394), between the Hawaiian Islands and the Philippines. It is conspicuously absent except at three stations along the line from Guam to Yokohama, being replaced at corresponding depths by volcanic mud. It is probable that this belt of volcanic mud does not extend far from the range of volcanic islands along which the cable route passes.

The least depth at which a distinctly red clay deposit has been noted is at station 680—2,010 fathoms. It is always found in abyssal depths. Ordinarily, as the contour line rises above the 2,500-fathom mark foraminifera rapidly increase in numbers and perfection of form, and soon justify the classification of the deposit under the head of globigerina ooze.

Globigerina ooze.—*Globigerina* ooze is defined as a deposit containing over 30 per cent of calcium carbonate, principally in the form of minute shells of foraminifera. Other organic remains commonly found in this deposit are sponge spicules, radiolaria, diatoms, and the very minute coccoliths and rhabdoliths. As a rule, in this part of the Pacific Ocean globigerina ooze will be found wherever the depth is less than 2,200 fathoms. The exceptions are found in the region of volcanic islands or submarine volcanic peaks where the foraminifera seem to be overwhelmed by volcanic sand, and in the vicinity of island shores where coral sand or blue or green mud may predominate. The globigerina ooze, wherever found on the line of this survey, is composed principally of the few species (about 20) of foraminifera known to be pelagic. Bottom living species are rare and individually few in number. The proportion of mineral matter, other than calcium carbonate, in this deposit is relatively small. Manganese concretions are generally present and sometimes quite numerous, and fragments of pumice are common. Crystals and spherules of phillipsite are often noted. The finer mineral fragments are quite lost in the mass of foraminifera, but appear when the latter are dissolved out with acid. At one station—643, 1,757 fathoms—the cavities of very many of the shells were found to be filled with a siliceous deposit forming complete casts of the

interior of the shells, even to the minute foramina. These casts are also noted twice in volcanic mud (stations 991 and 1065). Doubtless examples of these casts might be found in many other samples of globigerina ooze.

Diatom ooze.—Diatom ooze is the name given to a deep-sea deposit of which the principal constituent is the siliceous frustules of diatoms. Previous to this survey such a deposit had not been found in any tropical waters, and was supposed to be "confined to the Southern or Antarctic oceans, or to the northern parts of the North Pacific." Unexpectedly, therefore, many distinct patches of characteristic diatom ooze were found on the line, especially between Guam and Luzon, latitude $14^{\circ} 28'$ to $14^{\circ} 50'$ north, and longitude 136° to $130^{\circ} 30'$ east. Along this tract, about 300 miles in length, diatom ooze was recovered at stations, as follows:

Diatom ooze.

Station.	Latitude.			Longitude.			Depth.
	°	'	"	°	'	"	
743.....	14	28	00	136	00	00	3, 118
744.....	14	26	30	135	50	30	2, 879
746.....	14	24	00	135	31	00	2, 788
747.....	14	23	00	135	21	00	2, 731
750.....	14	25	00	134	51	30	2, 679
752.....	14	26	00	134	34	00	2, 432
764.....	14	29	00	133	56	15	2, 487
776.....	14	43	30	131	55	45	3, 283
781.....	14	48	30	131	03	00	3, 252
784.....	14	50	00	130	42	00	3, 547
920.....	14	31	15	132	42	30	3, 327
939.....	14	37	40	136	00	00	2, 838
959.....	14	13	00	139	34	00	3, 042

Between Guam and Midway Islands diatom ooze of the same nature appears at stations 559, 1710, and 1724. Also at stations 314 and 350 broken frustules of *Coscinodiscus rex* are noted.

As may be seen from the above table, the depths varied from 2,432 to 3,658 fathoms. In appearance the typical examples are greyish-white in color, shading off to a pale yellowish-brown wherever the fine red clay mud is present in any considerable proportion. In consistence it is mucilaginous, but is readily disintegrated by shaking with water. Radiolaria are generally rather numerous in this deposit. Mineral fragments are few. In all the specimens examined the diatoms belong almost exclusively to a single species identified by Professor Mann as *Coscinodiscus rex* Wallich. This is one of the largest diatoms known, having a diameter of about 0.8 millimeter, and is plainly visible to the naked eye. In form it resembles a minute pill box, with slightly rounded corners. The two valves (bottom and cover) are held together by a broad circumferential band. The valves are extremely thin and fragile, and the markings exceedingly delicate.

In some instances complete frustules are found, but usually the valves are separated and often much broken. A peculiar feature of this deposit is the strict limitation of the patches. Nearly pure diatom ooze may be recovered from one station, and at the next, five miles away, not a diatom appear in the deposit.

Radiolarian ooze.—No well-marked example of radiolarian ooze has been found in the specimens examined. Though radiolaria are noted in most of the samples, nowhere do they appear as a dominating constituent of the deposit. They are most numerous in the diatom oozes, where they are generally conspicuous by the number of individuals, but the number of species represented is not great.

Volcanic mud.—This is a deposit found in the neighborhood of volcanic islands or submarine volcanic peaks. Its characteristic constituents are pumice, glass, ashes, and the débris of volcanic rocks. It is often mixed with a considerable proportion of foraminifera when taken from depths less than 2,000 fathoms. Most often it is dark gray in color, and is readily disintegrated by shaking with water, being devoid of the sticky quality of red clay. This deposit is noted about the islands of Oahu and Guam, and nearly the whole distance from Guam to Yokohama, where the route passes along nearly parallel to the Ladrone and Bonin groups of volcanic islands, and at no great distance therefrom. The most conspicuous mineral constituent of this deposit is volcanic glass. It appears in various forms, the most frequent being the fibrous or filamentous variety. This has the appearance of having been drawn out when in a plastic state, sometimes into long, extremely fine threads, more commonly into larger threads or ribands, furrowed longitudinally, broken into short pieces, and always colorless and transparent as the finest artificial product. Another form is more massive, ragged in outline, dark brown, translucent, with numerous large, rounded cavities, and not so conspicuously suggestive of having been drawn out while cooling. A third variety consists of very fine, angular, perfectly transparent and colorless fragments, which often make up the bulk of the washed sediment. Red palagonite, coating fragments of other minerals is more frequently present in this deposit than in any other.

Blue mud.—Blue mud is the deposit generally found in inclosed or partially inclosed seas, and in the waters bordering continental land. It is composed for the most part of the débris carried out from the land by rivers or other currents. The few specimens collected by the *Nero* are blue-black in color, on the sides of the vial exposed to the light of a dark steel-blue with metallic luster, and iridescent. The color is said to be due to the presence of organic matter and iron sulphide. The odor of hydrogen sulphide is evident in all the well-corked vials of this mud. Except in deep waters foraminifera are more or less numerous. Radiolaria and diatoms are generally present, sometimes

in large numbers. Blue mud appears on the line of this survey only off the coasts of Luzon and Japan.

Green mud.—Green mud is found under the same conditions as blue mud. It is said to owe its color generally to the presence of the olive-green mineral glauconite, but sometimes to the presence of organic matter and its reducing action upon iron peroxide. In some instances the green color of the specimens has turned a bluish-black since recovery, and from present appearances would be called blue mud. In all the specimens of green mud the tinge of green is faint, and the greenish grains of sand comprise but a small part of the sediment. A large part of the coloration must be due to extremely minute amorphous mineral matter, since the supernatant water in the settling-glass remains cloudy and tinged with green after standing for an hour, and is not cleared or decolorized by nitro-hydrochloric acid. No glauconitic casts of foraminifera have been noted in these specimens. Green mud is recorded at several stations in Dingala Bay, coast of Luzon, and at all stations but one from No. 1217 to the anchorage near Yokohama, a distance of about 70 miles.

RECORD OF THE DETAILED EXAMINATION OF SELECTED SPECIMENS OF DEPOSITS FROM STATIONS ON THE OUTWARD VOYAGE OF THE NERO.

(A) HONOLULU TO MIDWAY ISLANDS.

Station 1.—923 fathoms. Volcanic mud. Sediment, after removal of "fine washings" by decantation, contains many foraminifera, a few sponge spicules, radiolarians, and diatoms. About 30 per cent of the sediment consists of fragments of volcanic rock and pumice. Many minute magnetic particles.

Station 4.—1,393 fathoms. Volcanic mud. Foraminifera numerous; sponge spicules, radiolarians, and diatoms few. Fine volcanic sand in small proportion.

Station 6.—2,438 fathoms. Volcanic mud. Foraminifera, radiolaria, diatoms, sponge spicules. Very fine volcanic ashes.

Station 11.—1,983 fathoms. Volcanic mud. Foraminifera (*Globigerina*, *Pulvinulina*, *Virgulina*, *Nonionina*, *Nodosaria*, *Hastigerina*). Radiolaria few. Diatoms few. About one-third the sediment fine volcanic sand.

Station 16.—2,438 fathoms. Volcanic mud. Color, pale yellowish brown. No foraminifera, a few radiolarians and diatoms. Mineral matter, fine volcanic sand. Many small fragments of pumice with minute manganese-iron concretions forming upon the surface.

Station 22.—2,673 fathoms. Red clay. No foraminifera; a few large radiolarians (*Oroplegma diplosphaera* Hæckel), mostly in fragments. Mineral fragments very small.

Station 28.—2,650 fathoms. Red clay. Fawn colored. No organic remains except a few radiolaria. Specimen consists almost entirely of fine amorphous clayey matter.

Station 36.—2,432 fathoms. Red clay. No foraminifera or radiolaria. Sediment, after removal of fine washings, small in quantity and composed entirely of minute particles of sand.

Station 46.—2,723 fathoms. Red clay. Fawn colored. Fine mud, with a few minute mineral fragments, none larger than 0.08 millimeter.

Station 65.—2,750 fathoms. Red clay. No organic remains except an occasional radiolarian. Mineral sediment small in quantity and exceedingly fine.

Station 81.—2,908 fathoms. Red clay. Mostly "fine washings;" a few minute radiolaria and mineral particles. No calcareous organisms.

Station 93.—1,463 fathoms. Globigerina ooze. Light grayish-brown. Broken shells of foraminifera; few complete ones. No coccoliths. Nodules of manganese; many rather coarse mineral fragments.

Station 100.—2,552 fathoms. Red clay. Fawn colored. Foraminifera few and much broken; no other organic remains. Coarse volcanic sand in large proportion.

Station 106.—2,002 fathoms. Specimen consists of three manganese-iron nodules, the largest about 12 millimeters in diameter. This is as large an object as the opening in the sounding cup would admit. The finer material was washed out of the cup during its return to the surface, the closure of the valve having been prevented by the nodules.

Station 110.—2,655 fathoms. Red clay. A few foraminifera. No other organic remains. Very small mineral sediment, principally volcanic glass.

Station 124.—1,726 fathoms. Globigerina ooze. Color, grayish-white. Sediment almost exclusively composed of foraminifera: *Orbulina*, *Globigerina*, *Pulvinulina*, *Polystomella*, *Verneuilina*, *Erhenbergina (hystrix)*, the latter rather frequent. Few mineral particles. A few coccoliths and rhabdoliths.

Station 125.—2,230 fathoms. Globigerina ooze. Color, brownish-white. Foraminifera: *Globigerina*, *Pulvinulina*, *Rotalia*, *Ehrenbergina (hystrix)*. Coccoliths; no radiolaria or diatoms. Nodules of phillipsite; decomposed pumice, coarse and fine.

Station 126.—2,627 fathoms. Red clay. Although this station is only 5 miles distant from the last, the foraminifera have entirely disappeared, and the deposit shows only amorphous matter, an occasional radiolarian, and a few mineral fragments.

Station 152.—3,026 fathoms. Red clay. Only a few particles larger than 0.3 millimeter. A single fragment of an arenaceous foraminifera (*Psammosphæra fusca*). No calcareous organisms. Fragments of large radiolarian (*Oroplegma*). Minute manganese concretions. Fine sand.

Station 163.—2,603 fathoms. Red clay. Fawn colored. No foraminifera; many radiolaria; few diatoms; sponge spicules. Mineral fragments very small in size and quantity.

Station 165.—2,135 fathoms. Globigerina ooze. Color, pale yellowish-brown. Sediment principally pelagic foraminifera; many coccoliths. Few mineral fragments.

Station 166 to 174.—1,593 to 2,111 fathoms. Globigerina ooze. Color varies from nearly white to pale yellowish-brown, according to

the proportion of foraminifera, which latter seems to be intimately related to the depth. Foraminifera: *Globigerina*, *Orbulina*, *Hastigerina*, *Pulvinulina*, *Pullenia*, *Miliolina*, *Ehrenbergina*, *Cyclammina*, *Virgulina*, *Uvigerina*, *Lagena*, *Discorbina*, *Polystomella*, *Nodosaria*, *Sphaeroidina*. Coccoliths more or less numerous, rhabdoliths few; sponge spicules; radiolaria not numerous except at station 174; diatoms few. Mineral fragments very few.

Station 175.—1,239 fathoms. This specimen vial contained only a few brownish-black fragments of a manganese nodule.

Station 185.—2,757 fathoms. Red clay. Brown-gray. Very fine mud, with a few sponge spicules, radiolaria, and an occasional diatom.

Station 187.—2,473 fathoms. Globigerina ooze. Color, light gray. The washed sediment consists of broken foraminifera, radiolaria, diatoms, and a very little fine sand.

Station 189.—1,813 fathoms. Globigerina ooze. Grayish-white. Foraminifera: *Biloculina*, *Orbulina*, *Pulvinulina*, *Uvigerina*, *Globigerina*, *Nodosaria*, *Lagena*, *Pullenia*, *Virgulina*, *Polystomella*. Coccoliths and rhabdoliths not numerous; occasional small radiolaria and diatoms. Mineral fragments very few.

(B) MIDWAY ISLANDS TO GUAM.

Station 205.—2,167 fathoms. Globigerina ooze. Light brown. Foraminifera mostly in fragments. A few radiolaria; many coccoliths. Mineral particles rare.

Station 209.—82 fathoms. Coral sand. Fragments of coral rock. Foraminifera (*Amphistigina*), polyzoa, and univalve mollusks. (This is the only specimen from a sounding less than 100 fathoms.)

Station 211.—2,322 fathoms. Red clay. Color, light brown. Specimen consists almost exclusively of fine washings. A few broken foraminifera, an occasional radiolarian, and the usual mineral fragments.

Station 225.—2,926 fathoms. Red clay. Total sediment consists of fine washings, with an occasional radiolarian and sponge spicule and a few small fragments of volcanic glass.

Station 238.—3,012 fathoms. Red clay. No effervescence with acid. No organic remains, except rarely a sponge spicule or fragments of a radiolarian. The few mineral particles are minute, colorless, transparent, vitreous fragments.

Station 248.—3,168 fathoms. Red clay. Light brown. A few radiolaria; no other organic remains. No effervescence with acid. Mineral particles very small, transparent fragments.

Station 257.—3,250 fathoms. Red clay. No calcareous organisms; a few radiolaria and sponge spicules. A large sediment of mineral fragments in great variety. Numerous small manganese nodules. Crystals and spherules of phillipsite.

Station 271.—3,240 fathoms. Red clay. Light brown, extremely fine mud. An occasional radiolarian; no other organic remains. No mineral particles larger than 0.08 millimeter.

Station 285.—3,089 fathoms. Red clay. Many minute manganese concretions.

Station 295.—3,274 fathoms. Red clay. Many small concretions of manganese and crystals of phillipsite, single and crossed.

Station 314.—3,237 fathoms. Red clay. Extremely fine mud. No mineral particles exceeding 0.08 millimeter in diameter. Gelatinous masses containing great numbers of fragments of large diatoms. (*Coscinodiscus rex* Wallich.) A few radiolaria.

Station 331.—2,997 fathoms. Red clay. Color, brown. No calcareous organisms. At least one-half of the washed sediment consists of crystals of phillipsite. Many small nodules of manganese. This specimen is unique in the preponderance of clear-cut crystals of phillipsite.

Station 335.—2,845 fathoms. Red clay. Light brown, very fine mud. No calcium carbonate. A few radiolarians. Minute nodules of manganese; a few crystals of phillipsite and glassy mineral fragments.

Station 336.—2,424 fathoms. Red clay. Broken shells of foraminifera begin to appear. Active effervescence with acid. Distance from preceding station about 11 miles; difference in depth, 421 fathoms.

Station 338.—2,128 fathoms. Globigerina ooze. Fawn color. Contains a large proportion of foraminifera, mostly broken and corroded; a few coccoliths. Fragments of pumice; many concretions of manganese of considerable size (6 millimeters), spherules of phillipsite, and minute glassy fragments.

Station 345.—1,173 fathoms. Pure globigerina ooze. Color, white, with slight shade of brown. Sediment composed almost entirely of perfect foraminifera (*Globigerina*, *Pulvinulina*, and *Orbulina*), with rarely a grain of sand.

Station 350.—2,240 fathoms. Red clay. Reddish-brown mud, containing many foraminifera, diatoms (fragments of *Coscinodiscus*), manganese concretions of considerable size (6 millimeters), crystals and spherules of phillipsite, and volcanic mineral particles.

Station 352.—2,568 fathoms. Red clay. Only an occasional corroded fragment of a foraminifer. The usual small manganese nodules, crystals of phillipsite, and other minerals.

Station 356.—2,897 fathoms. Red clay. Washed sediment very fine, consisting of a few minute manganese concretions and an abundance of single and crossed crystals of phillipsite.

Station 361.—2,268 fathoms. Red clay. Fawn-colored mud. Washed sediment composed largely of manganese nodules of consid-

erable size (up to 6 millimeters), and a small proportion of foraminifera mostly broken and corroded.

Station 362.—1,937 fathoms. Typical globigerina ooze. Pinkish-white. Very little fine washings. Sediment almost entirely foraminifera (*Globigerina*, *Orbulina*, *Pulvinulina*, *Lagena*, *Cassidulina*, *Ehrenbergina*, *Virgulina*, *Pullenia*). No radiolaria or diatoms observed. Very few coccoliths.

Station 369.—966 fathoms. Globigerina ooze. Foraminifera of the common pelagic species.

Station 373.—2,153 fathoms. Red clay. Fine yellowish-brown mud. A few foraminifera, mostly in corroded fragments. Manganese concretions, crystals and spherules of phillipsite, and minute magnetic particles.

Station 376.—2,780 fathoms. Red clay. Yellowish-brown. Almost entirely fine washings. No organic remains; very few mineral fragments.

Station 385.—720 fathoms. Globigerina ooze. (*Globigerina*, *Orbulina*, *Pulvinulina*, *Sphaeroidina*, *Cristellaria*.) An occasional radiolarian and bit of sand.

Station 390.—3,006 fathoms. Red clay. Exceedingly fine brown mud. Only separable residue a few minute manganese concretions, crystals of phillipsite, and discoid radiolaria.

Station 400.—3,159 fathoms. Red clay. Light yellowish-brown, very fine mud, containing no organic remains, very few mineral fragments exceeding 0.08 millimeter in diameter, and many minute crystals of phillipsite.

Station 411.—3,188 fathoms. Red clay. Very fine brown mud. A few radiolarians and diatoms, small crystals of phillipsite, and minute glassy mineral fragments.

Station 427.—1,997 fathoms. Globigerina ooze. Grayish-white. Foraminifera much broken. A few perfect specimens of *Ehrenbergina hystrix*. Rather large manganese concretions, fragments of pumice and spherules of phillipsite. Minute fragments of volcanic glass.

Station 451.—3,150 fathoms. Red clay. Yellowish-brown fine mud, containing a few radiolaria. The washed sediment consists of fine volcanic glass, and other mineral fragments varying in color from dark red-brown to light brownish-yellow. Minute manganese particles and a few crystals of phillipsite.

Station 460.—689 fathoms. Globigerina ooze. The usual pelagic foraminifera, a very few coccoliths and rhabdolites. Rarely a mineral fragment.

Station 463.—1,913 fathoms. Globigerina ooze. Specimen consists of a little globigerina ooze, and the fragments of a manganese

nodule, originally about 25 millimeters in diameter, probably broken by concussion of the sinker. Nucleus of the nodule is a porous, straw-colored fragment of pumice.

Station 478.—2,708 fathoms. Red clay. Dark brown. No foraminifera. A few radiolaria. Washed sediment mostly volcanic glass.

Station 498.—3,185 fathoms. Red clay. Numerous manganese concretions, crystals and nodules of phillipsite. No organic remains.

Station 506.—2,169 fathoms. Globigerina ooze. Color, brownish-white. Shells much broken. Foraminifera mostly pelagic; individual specimen of *Lagena gracilis*. A few coccoliths and rhabdoliths. Several rather large manganese nodules, 10 millimeters in diameter.

Station 521.—3,356 fathoms. Red clay. Chocolate color. No organic remains. Washed sediment consists of fine sand containing small manganese nodules, aggregated crystals of phillipsite, volcanic glass, and other minerals.

Station 530.—3,118 fathoms. Red clay. Brown mud. No organisms. Nodules of manganese and of phillipsite; decomposing pumice.

Station 541.—1,846 fathoms. Globigerina ooze. Creamy white. Foraminifera mostly pelagic. *Nonionina* noted. Very few mineral particles.

Station 559.—3,658 fathoms. Diatom ooze. Fine, light-brown mud. Relatively small quantity of clayey matter. Sediment composed of fragments of large diatoms; (*Coscinodiscus rex* Wallich) radiolaria in abundance. Volcanic ashes.

Station 575.—4,563 fathoms. Red clay. A fine, brown mud with a large percentage of clayey matter, and notable for the absence of manganese concretions. Washed sediment principally clear, transparent fragments of volcanic glass.

Station 591.—4,204 fathoms. Red clay. No organic remains. Mineral matter rather coarse volcanic fragments. Very few manganese concretions.

Station 600.—2,536 fathoms. Volcanic mud. Grayish-brown. No foraminifera; no effervescence with acid; a few radiolaria. Sediment consists of volcanic débris, with very little fine washings. Numerous manganese concretions, yellow-brown to red-brown particles of palagonite, and vitreous fragments.

Station 603.—1,745 fathoms. Volcanic mud. Gray, granular mud. About 25 per cent of the sediment consists of pelagic foraminifera; the remainder is a rather fine volcanic sand containing manganese concretions, palagonite, and vitreous fragments in large proportions. Many magnetic particles.

Station 613.—1,072 fathoms. Pure globigerina ooze. Pelagic foraminifera with few exceptions. One *Lagena globosa* and one *Gaudryina pupoides* noted.

Station 614.—3,230 fathoms. Red clay. No organic remains. Sediment of decomposing pumice and minute irregular fragments of perfectly transparent rock.

Station 615.—3,178 fathoms. Red clay. No organisms. Fine mineral fragments and occasional minute manganese nodules.

Station 637.—2,352 fathoms. Volcanic mud. Gray, granular non-coherent mud. Very little fine washings. Washed sediment contains a small proportion of foraminifera, arenaceous (*Jaculella*) and cretaceous, very many manganese nodules, volcanic glass, and other minerals not identified.

Station 643.—1,757 fathoms. Globigerina ooze. Brownish-gray. But little amorphous matter. Sediment mostly the usual pelagic forms of foraminifera, and fine, glassy mineral fragments. After action of acid there remain large numbers of white silicious casts of foraminifera, often quite perfect, even of the minute foramina of the shells.

Station 647.—605 fathoms. Globigerina ooze. Mostly pelagic foraminifera, with fine coral sand.

(C) GUAM TO LUZON.

Station 663.—457 fathoms. Coral sand. Blue-black (probably from chemical changes since collection). Contains shells of small univalve and bivalve mollusks, fragments of coral, foraminifera (*Pulvinulina*, *Cristellaria*, *Cassidulina*, *Miliolina*, *Nonionina*, *Amphistegina*, *Lagena*), manganese nodules, and much fine mineral sand.

Station 670.—1,376 fathoms. Volcanic mud. Yellowish-brown to black, very irregular fragments of transparent or translucent volcanic glass. The fragments have a resinoid luster, are porous, sharply angular, often fibrous, as if drawn out when in a semifluid state. A few foraminifera.

Station 674.—1,946 fathoms. Volcanic mud. Very pale yellowish-brown. Sediment consists of a few foraminifera and radiolaria, and a large proportion of minute splinters of volcanic glass. Many particles of palagonite.

Station 688.—1,346 fathoms. Globigerina ooze. Contains a very large number of manganese nodules.

Station 705.—2,710 fathoms. Red clay. No foraminifera; a few radiolaria and manganese nodules. Volcanic ashes.

Station 715.—2,639 fathoms. Red clay. Many manganese concretions coating fragments of volcanic minerals.

Station 722.—2,476 fathoms. Red clay. A chocolate-colored, very sticky mud. No organic remains. Granular coatings of manganese upon fragments of pumice and lumps of clay. Volcanic ashes.

Station 730.—2,761 fathoms. Red clay. Pale yellowish-brown, very fine mud. Rarely a radiolarian or diatom. Mineral matter small in quantity and minute in size

Station 740.—2,735 fathoms. Red clay. Yellowish-brown. No organisms noted. Many small manganese nodules; very little other mineral matter.

Station 743.—3,118 fathoms. Diatom ooze. A grayish-white mucilaginous mass, composed almost entirely of the more or less broken frustules of large diatoms, *Coscinodiscus rex* Wallich (identification by Prof. Albert Mann). Many radiolaria are found among the diatoms. There is very little clayey matter and few mineral fragments.

Station 744.—2,879 fathoms. Diatom ooze. Like the preceding specimen, except that it contains more clay, and radiolaria more numerous.

Station 745.—2,617 fathoms. Red clay. Extremely fine chocolate-colored mud with a few minute mineral fragments, but no diatoms or other organisms.

Station 746.—2,788 fathoms. About 9 miles from station 745. Diatom ooze. Same as station 743. Quite a large proportion of the valves in this specimen are unbroken. A few entire frustules.

Station 747.—2,731 fathoms. Ten miles from station 746. Red clay. A fine, sticky, deep yellowish-brown mud without trace of a diatom or other organism.

Station 748.—2,891 fathoms. Red clay. Same as station 747.

Station 749.—2,819 fathoms. Diatom ooze. Same as stations 743 and 746. The frustules are much broken, but belong to the same species, *Coscinodiscus rex*.

Station 750.—2,679 fathoms. Diatom ooze. Characters same as above. Many unbroken valves, and occasionally a complete frustule.

Station 751.—2,679 fathoms. Red clay. Dark yellowish-brown mud. No diatoms or radiolaria.

Station 752.—2,432 fathoms. Diatom ooze. Identical with station 743, except that the color is a darker gray.

Station 753.—1,913 fathoms. Globigerina ooze. Nearly the whole sediment consists of pelagic species of foraminifera; rarely a radiolarian; not a fragment of a *Coscinodiscus*.

Station 760.—1,560 fathoms. Globigerina ooze. Very few coccoliths. Foraminifera usual pelagic species, and *Nodosaria*, *Lagena* (*sulcata*), *Pulvinulina* (*pauperata*).

Station 764.—2,487 fathoms. Diatom ooze. Light yellowish-brown. Washed sediment consists of fragments of *Coscinodiscus rex* Wallich, with many radiolaria.

Station 770.—2,888 fathoms. Red clay. Very fine chocolate-colored mud. No organisms. Few minute mineral fragments.

Station 776.—2,383 fathoms. Diatom ooze. Pale yellowish-brown. Consists of diatoms (*Coscinodiscus rex*) with a considerable proportion of fine clay.

Station 777.—3,421 fathoms. Red clay. Color, brown. No diatoms; a few radiolaria. The usual minute manganese particles and fine mineral sand.

Station 781.—3,252 fathoms. Diatom ooze. Whole deposit consists of broken frustules of *Coscinodiscus*.

Station 783.—3,264 fathoms. Red clay. A single small manganese nodule and one arenaceous foraminifer (*Reophax*) noted. Residue, fine mud with minute vitreous fragments.

Station 784.—3,547 fathoms. Diatom ooze. Fine clayey matter predominates, but fragments of *Coscinodiscus* make a large proportion of the deposit.

Station 790.—3,119 fathoms. Red clay. Very fine yellowish-brown mud containing a few radiolaria and fine mineral particles.

Station 796.—2,670 fathoms. Red clay. Very fine mud, light chocolate color. Contains a few radiolaria, and mineral fragments rarely exceeding 0.08 millimeter in diameter.

Station 801.—3,298 fathoms. Red clay. Grayish-brown, not very adhesive mud, containing a few radiolaria and sponge spicules and a large proportion of very fine rock fragments.

Station 808.—2,855 fathoms. Red clay. Very fine light-brown mud. A few radiolaria and sponge spicules and a small proportion of mineral fragments.

Station 812.—3,130 fathoms. Red clay. Very fine yellowish-brown mud, leaving, after washing, a small sediment of radiolaria and fine mineral particles.

Station 818.—3,182 fathoms. Red clay. Brown mud containing fine sand and a few radiolaria and sponge spicules.

Station 822.—2,427 fathoms. Red clay. Dark brown. No organic remains. Washed sediment mostly colorless transparent mineral fragments and fibrous volcanic glass.

Station 828.—1,390 fathoms. Blue mud. Had distinct odor of hydrogen sulphide when vial was first opened. Brownish-gray color. Contains a few foraminifera, radiolaria, and casts. Much the largest part of the sediment consists of angular transparent fragments of rock, for the most part less than 0.08 millimeter diameter.

Station 833.—2,740 fathoms. Red clay. Light gray brown very fine mud. No effervescence with acid. Radiolaria, sponge spicules and a few diatoms. Minute angular rock fragments in large proportion.

Station 850.—157 fathoms. Green mud. Dark greenish brown. A few sponge spicules; no foraminifera or radiolaria noted. Washed sediment consists of angular rock fragments, many of them various shades of green.

(D) GUAM TO YOKOHAMA, JAPAN.

Station 990.—859 fathoms. Coral sand. Color, gray. Fragments of coral rock. Many foraminifera; a few radiolaria. Many small manganese concretions and particles of palagonite, the latter being unusually numerous. Mineral fragments in quantity, angular, many of them green.

Station 995.—2,091 fathoms. Volcanic mud. No calcium carbonate. No organic remains. Very little fine washings. Sediment principally volcanic glass.

Station 1000.—1,947 fathoms. Volcanic mud. Very little fine mud. An occasional foraminifer and radiolarian. Sediment mostly fibrous volcanic glass.

Station 1006.—1,847 fathoms. Volcanic mud. Brownish gray. Foraminifera few; radiolaria rather numerous. Mineral matter fine volcanic glass.

Station 1010.—2,082 fathoms. Volcanic mud. Few foraminifera. Sediment, fine angular particles of volcanic sand. Very few of the fibrous fragments of glass so plentiful at stations 1000 and 1006.

Station 1016.—2,375 fathoms. Volcanic mud. Color, dark brown. About 50 per cent of fine washings; few foraminifera. The remainder consists of fine angular particles of volcanic sand.

Station 1026.—2,025 fathoms. Volcanic mud. Grayish brown. Few foraminifera; radiolaria rather numerous. Sediment chiefly angular mineral fragments in great variety. Fine washings 35 per cent of total sediment, but a large proportion of these washings consists of minute fragments of minerals.

Station 1036.—2,155 fathoms. Volcanic mud. Light brown, finely granular, nonadhesive mud, containing a few foraminifera and a relatively small amount of amorphous matter. The remainder is made up of fine angular mineral fragments.

Station 1045.—2,330 fathoms. Volcanic mud. Dark brown. No foraminifera, a few radiolaria, about 25 per cent of amorphous matter and volcanic sand.

Station 1055.—2,028 fathoms. Volcanic mud. Dark brown. No foraminifera or diatoms, radiolaria rather numerous. Washed sediment consists of manganese concretions and angular, colorless, transparent mineral fragments; many palagonite particles.

Station 1065.—1,321 fathoms. Volcanic mud. Light gray, granular, nonadhesive. Many foraminifera and siliceous casts; occasional radiolaria; much fine volcanic sand in angular particles.

Station 1074.—483 fathoms. Volcanic sand. Specimen consists of comparatively coarse volcanic sand, with a few foraminifera.

Station 1084.—2,313 fathoms. Volcanic mud. Light brownish gray, granular. An occasional foraminifer; many radiolaria. Much

volcanic glass, some of it of the brown porous variety, some filamentous, and the remainder sharp, angular, perfectly transparent fragments.

Station 1094.—3,495 fathoms. Red clay. Brown, sticky mud, consisting largely of amorphous clayey matter, with a small quantity of mineral fragments of a distinctly volcanic character.

Station 1104.—2,214 fathoms. Volcanic mud. Specimen consists of a single lapillus of brown porous volcanic glass about 6 millimeters in diameter.

Station 1110.—2,870 fathoms. Volcanic mud. A few arenaceous foraminifera (*Rhabdammina*, *Haplophragmium*) and radiolaria. Sediment composed almost entirely of volcanic glass.

Station 1120.—1,710 fathoms. Volcanic mud. Yellowish-brown granular mud, containing a few foraminifera, many radiolaria, and much volcanic sand, of which the larger particles are dark-brown glass.

Station 1126.—927 fathoms. Volcanic mud. A few foraminifera (*Globigerina*, *Pulvinulina*, *Pullenia*, *Uvigerina*). The rest of sediment volcanic sand.

Station 1132.—2,950 fathoms. Volcanic mud. Brownish gray, granular. No foraminifera; few radiolaria. Large proportion of volcanic sand, principally brown glass, and olive-green rounded mineral fragments.

Station 1142.—2,682 fathoms. Volcanic mud. No effervescence with acid. Many radiolaria; a few diatoms. Small manganese concretions; lapilli and fine fragments of volcanic glass.

Station 1151.—1,686 fathoms. Globigerina ooze. Very light gray. Contains 30 per cent or more of foraminifera, coccoliths, and rhabdoliths. Small manganese concretions and vitreous mineral fragments, with many red particles of palagonite.

Station 1168.—2,933 fathoms. Volcanic mud. No foraminifera, a few radiolaria and diatoms; large proportion of rather coarse sand and fine volcanic glass.

Station 1185.—1,491 fathoms. Volcanic mud. Color, light gray, slowly turning black with time. Many foraminifera (*Globigerina*, *Orbulina*, *Pullenia*, *Polystomella*, *Biloculina*, *Nonionina*, *Nodosaria*); radiolaria numerous; diatoms few. Many manganese concretions; much colorless volcanic glass, palagonite, and a variety of unidentified minerals.

Station 1197.—1,698 fathoms. Volcanic mud. Light gray, becoming black. A few foraminifera; very many radiolaria and diatoms. Manganese concretions, volcanic glass, palagonite, and various unidentified mineral fragments.

Station 1207.—665 fathoms. Blue mud. Blue black. Distinct odor of hydrogen sulphide, increased by addition of hydrochloric acid.

Contains a few small foraminifera and radiolaria. Coarse mineral fragments, many of them black. Many fragments coated with red palagonite.

Station 1217.—934 fathoms. Green mud. Dark gray. Has evidently changed color since collection, for it is noted on record as "gr. m.," green mud. Marked odor of hydrogen sulphide. The washed sand consists principally of vitreous fragments, some of them dark brown and nearly opaque, others clear and transparent. Occasional pale-green grains. No casts.

Station 1237.—613 fathoms. Green mud. Turned black from development of hydrogen sulphide since collection. A few foraminifera; very many diatoms; no radiolaria. Much fine sand.

ABSTRACT OF THE OFFICIAL RECORD OF SOUNDINGS.

HAWAIIAN ISLANDS TO MIDWAY ISLANDS.

[Columns marked "Deposit" and "Remarks" supplied by the compiler.]

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
1	May 6	21 12 00	158 11 00	923	78			gn. m. fn. s.	Volcanic mud.	
2	do	21 12 00	158 11 00	904	78			gn. m. fn. s.		No specimen.
3	do	21 25 00	158 28 00	1,299	81	78				Do.
4	do	21 30 00	158 30 00	1,393	80	78		br. m. fn. s.	Volcanic mud.	
5	do	21 53 00	158 30 00	964	73	76				Do.
6	do	22 04 00	158 30 00	2,201	74	76		fn. br. m.	Volcanic mud.	
7	May 7	22 12 00	158 40 00	2,032	74	74				Do.
8	do	22 14 00	158 42 00	1,802						Do.
9	do	22 21 00	158 51 00	2,242	74	74		fn. br. m.	Volcanic mud.	
10	do	22 23 00	158 54 00	2,098	74	75		fn. br. m.	do	
11	do	22 30 00	159 05 00	1,963	75	75		fn. br. m.	do	With forami- nifera.
12	do	22 33 00	159 07 00	1,924	74	75		fn. br. m.	do	
13	do	22 40 00	159 16 00	1,801	79	75				No specimen.
14	do	22 41 00	159 20 00	1,866	79	76				Do.
15	do	22 50 00	159 29 00	2,443	75	76		br. m. fn. s.	Volcanic mud.	
16	do	22 51 00	159 30 00	2,438	74	76		br. m. fn. s.	do	Fine volcanic sand.
17	do	22 58 00	159 37 00	2,709			35.1	br. m. fn. s.	do	
18	do	23 00 00	159 38 00	2,864	74	75				No specimen.
19	do	23 05 00	159 47 00	2,700	73	75		br. m.	Red clay	
20	do	23 07 00	159 50 00	2,704	74	75		br. m.	do	
21	May 8									No record.
22	do	23 13 00	159 59 00	2,673				br. m.	Red clay	
23	do	23 13 00	160 01 00	2,664	75	74	35.3	lt. br. m. dk. s.	do	
24	do	23 18 00	160 10 00	2,644	76	75		br. m.	do	
25	do	23 20 00	160 12 00	2,650	78	75		br. m.	do	
26	do	23 24 00	160 22 00	2,704	78	75		br. m. fn. s.	do	
27	do	23 25 00	160 23 00	2,788	78	75		br. m. fn. s.	do	
28	do	23 29 00	160 33 00	2,650	79	76		br. m. fn. s.	do	
29	do	23 29 00	160 35 00	2,652	79	76		br. m. fn. s.	do	
30	do	23 34 00	160 40 00	2,648	75	76		br. m. fn. s.	do	
31	May 9	23 35 00	160 48 00	2,724	75	74		br. m. fn. sp.	do	
32	do	23 39 00	161 00 00	2,699	75	75		br. m. fn. sp.	do	
33	do	23 40 00	161 03 00	2,572	74	75		br. m. fn. sp.	do	
34	do	23 45 00	161 15 00	2,466	76	75		br. m. fn. sp.	do	
35	do	23 46 00	161 18 00	2,467	75	75	35	br. m. fn. sp.	do	
36	do	23 51 00	161 31 00	2,432	75	74		br. m. fn. sp.	do	
37	do	23 52 00	161 33 00	2,453	79	74		br. m. fn. sp.	do	
38	do	23 56 00	161 45 00	2,471	78	76		br. m. fn. sp.	do	
39	do	23 58 00	161 47 00	2,474	77	76		br. m. fn. sp.	do	
40	do	24 01 00	161 56 00	2,435	75	76		br. m. fn. sp.	do	
41	do	24 02 00	161 58 00	2,484	75	76				No specimen.
42	do	24 06 00	161 08 00	2,574	75	76		br. m. fn. sp.	Red clay	
43	May 10	24 07 00	162 10 00	2,600	74	75		br. m. fn. sp.	do	
44	do	24 10 00	162 20 00	2,718	74	75		br. m. fn. sp.	do	
45	do	24 11 00	162 22 00	2,671	74	75				Do.
46	do	24 15 00	162 32 00	2,723	74	75		br. m.	Red clay	
47	do	24 16 00	162 35 00	2,706	79	75		br. m.	do	
48	do	24 20 00	162 47 00	2,722	79	75.5		br. n.	do	
49	do	24 21 00	162 49 00	2,726	81	76	35	br. m.	do	
50	do	24 23 00	163 01 00	2,732	81	75		br. m.	do	
51	do	24 24 00	163 04 00	2,739	79	75		br. m.	do	
52	do	24 26 00	163 15 00	2,742	79	75	35	br. m.	do	
53	do	24 26 00	163 18 00	2,759	77	75		br. m.	do	

Abstract of the official record of soundings—Continued.

HAWAIIAN ISLANDS TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
54	May 11	24 28 00	163 28 00	2,779	73	75	35	br. m.	Red clay...	
55do	24 28 00	163 30 00	2,765	73	75		br. m.do	
56do	24 30 00	163 41 00	2,742	76	75	35	br. m.do	
57do	24 31 00	163 44 00	2,727	77	75		br. m.do	
58do	24 32 00	163 54 00	2,722	80	76		br. m.do	
59do	23 33 00	163 57 00	2,718	79	75				No specimen.
60do	24 37 00	164 07 00	2,722	77	75	35	br. m.	Red clay...	
61do	24 38 00	164 09 00	2,723	75	75		br. m.do	
62do	24 41 00	164 18 00	2,737	75	75	35	br. m.do	
63do	24 42 00	164 20 00	2,746	74	74		br. m.do	
64	May 12	24 46 00	164 29 00	2,746	74	74	35	br. m.do	
65do	24 46 00	164 31 00	2,750	74	74		br. m.do	
66do	24 50 00	164 41 00	2,775	73	74	35	br. m.do	
67do	24 51 00	164 43 00	2,780	74	74		br. m.do	
68do									No record.
69do	24 55 00	164 52 00	2,772	75	75		br. m.	Red clay...	
70do	24 55 00	164 53 00	2,765	75	75				No specimen.
71do	24 59 00	165 04 00	2,751	75	76				Do.
72do	25 03 00	165 13 00	2,744	74	76				Do.
73do	25 04 00	165 15 00	2,715	73	75				Do.
74do	25 07 00	165 24 00	2,741	74	74		br. m.	Red clay...	
75do	25 08 00	165 26 00	2,741	74	74				Do.
76	May 13	25 10 00	165 34 00	2,725	74	75		br. m.	Red clay...	
77do	25 11 00	165 36 00	2,719	74	74	35	br. m.do	
78do	25 13 00	165 44 00	2,735	71	74	35	br. m.do	
79do	25 14 00	165 46 00	2,720	71	74		br. m.do	
80do	25 18 00	165 59 00	2,750	72	74	35.1	br. m.do	
81do	25 21 00	166 11 00	2,908	78	74.5		br. m.do	
82do	25 22 00	166 13 00	2,760	75	73		br. m. fn. sp.do	
83do	25 25 00	166 23 00	2,758	73	74	35	br. m. fn. sp.do	
84do	25 25 00	166 25 00	2,754	73	74		br. m. fn. sp.do	
85do	25 28 00	166 37 00	2,756	73	73		br. m. fn. sp.do	
86	May 14	25 29 00	166 39 00	2,755	72	73		br. m. fn. sp.do	
87do	25 32 00	166 51 00	2,765	72	73	35	br. m. fn. sp.do	
88do	25 33 00	166 54 00	2,770	73	74		br. m. fn. and crs. sp.do	Coarse min- eral frag- ments.
89do	25 36 00	167 05 00	2,535	74	74	35	br. m. fn. and crs. sp.do	
90do	25 37 00	167 07 00	2,299	74	73		br. m. fn. and crs. sp.do	
91do	25 38 00	167 09 00	1,983	76	74		fn. wh. s.	Globigerina ooze.	
92do	25 39 00	167 12 00	2,004	73	75		fn. wh. s.do	
93do	25 41 00	167 18 00	1,463	74	74		fn. wh. s.do	
94do	25 42 00	167 21 00	1,851	73	74				No specimen.
95do									Bottom not reached.
96do	25 46 00	167 30 00	2,269	73	74		br. m. wh. s. r.do	No specimen.
97do	25 52 00	167 42 00	2,114	73	74	35			Do.
98do	25 53 00	167 45 00	1,960	74	74		fn. wh. s.	Globigerina ooze.	
99do	25 56 00	167 52 00	1,895	73	74		rocky. G.		Do.
100	May 15	26 02 00	167 56 00	2,552	74	74		br. m. and G.	Red clay...	Coarse vol- canic sand.
101do	26 09 00	168 07 00	2,445	72	74		br. m. fn. sp.do	
102do	26 10 00	168 09 00	2,406	71	74		br. m. fn. sp.do	
103do	26 16 00	168 21 00	2,554	75	74	35	br. m. fn. sp.do	
104do	26 17 00	168 23 00	2,536	78	74		br. m. fn. sp.do	
105do	26 22 00	168 33 00	2,370	74	74		br. m. fn. sp.do	
106do	26 23 00	168 35 00	2,002	75	74		wh. s. R.	Globigerina ooze.	
107do	26 26 00	168 46 00	2,492	75	73	35.6	br. m. s. bk. sp.	Red clay...	
108do	26 27 00	168 48 00	2,527	75	73		br. m. s. bk. sp.do	
109do	26 29 00	168 57 00	2,662	74	74		br. and gr. m.do	
110do	26 30 00	168 59 00	2,655	73	73		br. and gr. m.do	
111do	26 32 00	169 08 00	2,642	73	73	35	br. m.do	
112	May 16	26 32 00	169 10 00	2,614	72	73		br. m. wh. sp.do	
113do	26 34 00	169 20 00	2,493	72	73	35	br. m. wh. sp.do	
114do	26 34 00	169 22 00	2,541	73	74		br. m.do	
115do	26 36 00	169 31 00	2,494	73	75	35	br. m.do	
116do	26 39 00	169 43 00	2,514	74	74		br. m.do	
117do	26 39 00	169 45 00	2,493	75	74		br. m.do	
118do	26 45 00	169 59 00	2,539	75	75	35.7	br. m.do	
119do	26 46 00	170 02 00	2,523	73	75		br. m.do	

Abstract of the official record of soundings—Continued.

HAWAIIAN ISLANDS TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	<i>Fath- oms.</i>	°	°	°			
120	May 16	26 49 00	170 13 00	2,534	74	74	35	br. m.	Red clay...	
121do....	26 52 00	170 25 00	2,562	73	75		br. m.do....	
122do....	26 55 00	170 37 00	2,571	72	74		br. m. fn. sp.do....	
123	May 17	26 56 00	170 40 00	2,568	71	74		br. m. fn. sp.do....	
124do....	26 59 00	170 52 00	1,726	71	74		wh. s. bk. sp.	Globigerina ooze.	
125do....	27 00 00	170 55 00	2,230	71	74		br. m. G.do....	Pumice.
126do....	27 02 00	171 01 00	2,627	71	74		br. m. fn. sp.	Red clay...	
127do....	27 03 00	171 08 00	2,636	74	75	35	br. m. fn. sp.do....	
128do....	27 11 00	171 19 00	2,675	76	74		br. m.do....	
129do....	27 12 00	171 21 00	2,678	76	75		br. m.do....	
130do....	27 17 00	171 36 00	2,706	77	74		br. m.do....	
131do....	27 22 00	171 49 00	2,716	72	74	44	br. m.do....	
132do....	27 26 00	172 00 00	2,732	72	74		br. m.do....	
133do....	27 26 00	172 02 00	2,734	72	74		br. m.do....	
134do....	27 29 00	172 11 00	2,734	70	74		br. m.do....	
135	May 18	27 30 00	172 13 00	2,734	70	74		br. m.do....	
136do....	27 32 00	172 22 00	2,749	70	74		br. m.do....	
137do....	27 33 00	172 24 00	2,812	70	74		br. m.do....	
138do....	27 36 00	172 32 00	2,788	71	75	51.7	br. m.do....	
139do....	27 36 00	172 34 00	2,773	72	75		br. m.do....	
140do....	27 39 00	172 44 00	2,763	71	75	44.8	br. m.do....	
141do....	27 43 00	172 55 00	2,801	71	75		br. m.do....	
142do....	27 47 00	173 05 00	2,865	73	75	do....	No specimen.
143do....	27 47 00	173 06 00	2,919	70	74		br. m.	Red clay...	
144do....	27 51 00	173 16 00	2,873	69	73	35	br. m.do....	
145do....	27 55 00	173 25 00	2,863	69	72	35	br. m.do....	
146	May 19	27 59 00	173 34 00	2,898	67	70		br. m.do....	
147do....	28 03 00	173 43 00	2,910	67	71	35	br. m.do....	
148do....	28 07 00	173 52 00	2,925	69	71		br. m.do....	
149do....	28 12 00	174 03 00	2,928	69	74	35	br. m.do....	
150do....	28 13 00	174 06 00	2,932	76	74		br. m.do....	
151do....	28 18 00	174 17 00	2,945	72	72		br. m.do....	
152do....	28 21 00	174 30 00	3,026	67	72	35	br. m.do....	
153do....	28 23 00	174 41 00	2,958	68	70		br. m.do....	
154do....	28 25 00	174 51 00	2,943	67	70		br. m.do....	
155do....	28 27 00	175 02 00	2,875	69	69		br. m.do....	
156	May 20	28 28 00	175 09 00	2,827	66	68		br. m.do....	
157do....	28 31 00	175 20 00	2,732	66	68	35	br. m.do....	
158do....	28 33 00	175 25 00	2,675	66	68		br. m.do....	
159do....	28 36 00	175 36 00	2,572	69	70		br. m.do....	
160do....	28 39 00	175 46 00	2,637	72	69		br. m.do....	
161do....	28 41 00	175 53 00	2,695	72	68		br. m. fn. sp.do....	
162do....	28 41 00	176 10 00	2,679	75	68		br. m. fn. sp.do....	
163do....	28 41 00	176 23 00	2,603	74	68	35	br. m. ers. sp.do....	
164do....	28 41 00	176 25 00	2,471	69	67		br. m. ers. sp.do....	
165do....	28 41 00	176 37 00	2,135	68	69		fn. co. s.	Globigerina ooze.	
166do....	28 41 00	176 40 00	1,850	69	71		fn. co. s.do....	
167do....	28 41 00	176 43 00	1,593	69	72		fn. co. s. and R.do....	Large mang. concretions.
168	May 21	28 41 00	176 45 00	1,667	68	72	do....	No specimen.
169do....	28 41 00	176 48 00	2,426	69	70	35do....	Do.
170do....	28 41 00	176 46 00	1,990	68	70		fn. and ers. co. s.	Globigerina ooze.	
171do....	28 39 00	176 46 00	1,913	68	70		fn. and ers. co. s.do....	
172do....	28 38 00	176 48 00	2,086	70	69		fn. and ers. co. s.do....	
173do....	28 43 00	176 39 00	2,111	71	72	35	fn. co. s. br. m.do....	
174do....	28 42 00	176 42 00	1,849	73	72		ers. co. s.do....	
175do....	28 41 00	176 45 00	1,239	73	72		blk. r. co.do....	Manganese nodule.
176do....	28 48 00	176 45 00	2,227	73	72	do....	No specimen.
177do....	28 50 00	176 49 00	2,633	71	72	do....	Do.
178do....	28 54 00	176 46 00	2,478	71	72		br. m. co. s. fn. sp.	Red clay...	
179do....	28 54 00	176 48 00	2,416	71	71		bk. r.do....	Manganese concretions.
180do....	28 54 00	176 50 00	2,893	70	69	do....	
181do....	28 51 00	176 56 00	2,836	69	68		br. m. fn. sp.	Red clay...	
182do....	28 47 00	177 01 00	2,865	69	70	35	br. m. fn. sp.do....	
183	May 22	28 44 00	177 07 00	2,796	69	70		br. m. fn. sp.do....	
184do....	28 41 00	177 12 00	2,805	68	70		br. m. fn. sp.do....	
185do....	28 39 00	177 15 00	2,757	69	70		br. m. fn. sp.do....	

Abstract of the official record of soundings—Continued.

HAWAIIAN ISLANDS TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur-face.	Bot-tom.			
	1899.	° / "	° / "	Fath-oms.	°	°	°			
186	May 22	28 36 00	177 20 00	2,539	69	70		bk. co. r.		No specimen.
187	do	28 32 00	177 22 00	2,473	71	70		co. s. and r.	Globigerina ooze.	
188	do	28 27 00	177 23 00	2,061	72	71		wh. co. s.	do	
189	do	28 25 00	177 24 00	1,813	72	71	35	wh. co. s.	do	
190	do	28 22 00	177 23 00	864	71	71		wh. co. s.	do	
191	do	28 20 00	177 24 00	51	71	70		co. co. s.	do	
192	do	28 20 00	177 23 00	155	71	70				
193	do	28 19 00	177 25 00	40	71	70				Do.
194	do	28 17 00	177 26 00	47	71	70				Do.
195	do	28 15 00	177 25 00	44	70	70				Midway Is-lands.

MIDWAY ISLANDS TO GUAM.

196	May 24	28 14 00	177 25 00	20				co. s.		No specimen. Do. Do. Do.
197	do	28 14 30	177 26 00	40				co. s.		
198	do	28 14 30	177 27 00	70				co. s.		
199	do	28 15 00	177 27 00	120				co. s.		
200	do	28 15 00	177 28 30	625	73	71		fn. wh. co. s.	Globigerina bl. sp. ooze.	
201	do	28 14 00	177 31 00	1,033	73	71		fn. wh. co. s.	do	
202	do	28 14 00	177 33 00	1,361	73	71		bl. sp.		
203	do	28 13 00	177 35 00	1,625	74	71	35	co. s. brk. sp.	do	
204	do	28 11 00	177 30 00	1,947	73	70		crs. co. s.	do	
205	do	28 07 00	177 42 45	2,167	73	70		fn. co. s. br.	do	
206	do	28 04 00	177 46 50	2,055	72	70		m.		
207	do	28 02 00	177 48 40	1,842	72	71		fn. co. s. br.	do	
208	do	27 58 00	177 51 40	1,351	72	71	35.2	fn. co. s.	do	
209	do	27 58 00	177 03 40	82	72	71		co.	Coral sand.	
210	do	27 57 00	177 41 15	1,718	72	71		crs. co. s.	Globigerina ooze.	
211	do	27 57 00	177 43 30	2,322	72	72		fn. co. s. br. m.	Red clay	A few foram-inifera.
212	do	27 52 00	177 44 00	2,036	72	72	35	crs. co. s.	Globigerina ooze.	
213	do	27 50 00	177 43 00	2,367	72	72				
214	do	27 46 00	177 41 00	2,539	72	72				
215	May 25	27 42 00	177 40 30	2,577	73	71		br. m.	Red clay	
216	do	27 39 00	177 41 00	2,592	73	72		br. m.	do	
217	do	27 35 00	177 42 00	2,619	72	72		br. m.	do	
218	do	27 29 00	177 46 00	2,632	73	72		br. m.	do	
219	do	27 26 00	177 52 00	2,621	73	72	35	br. m.	do	
220	do	27 22 00	178 05 30	2,654	73	72		br. m.	do	
221	do	27 19 00	178 16 30	2,768	73	73		br. m.	do	
222	do	27 16 00	178 29 00	2,850	73	73	35	br. m.	do	
223	do	27 13 00	178 40 30	2,884	74	75		br. m.	do	
224	do	27 10 00	178 51 30	2,905	75	75	35	br. m.	do	
225	do	27 09 00	179 01 15	2,926	73	74		br. m.	do	
226	May 26	27 08 00	179 10 30	2,939	75	75	35	br. m.	do	
227	do	27 08 00	179 12 30	2,934	74	75		br. m.	do	
228	do	27 07 00	179 21 30	2,934	74	75		br. m.	do	
229	do	27 06 00	179 23 30	2,934	75	75		br. m.	do	
230	do	27 06 00	179 32 45	2,956	75	75	35	br. m.	do	
231	do	27 35 00	179 42 30	2,948	76	74		br. m.	do	
232	do	27 03 00	179 53 15	2,960	80	75	39.4	br. m.	do	
233	do	27 01 00	East.	2,967	76	76	35	br. m.	do	
234	do	26 58 00	179 42 15	2,959	78	76		br. m.	do	
235	do	26 55 00	179 31 45	2,982	75	76	35	br. m.	do	
236	do	26 53 00	179 21 15	2,982	75	75		br. m.	do	
237	May 28	26 50 00	179 11 30	2,993	74	75	35	br. m.	do	
238	do	26 48 00	179 01 15	3,012	74	74		br. m.	do	
239	do	26 45 00	178 51 15	3,048	75	74	52	br. m.	do	
240	do	26 42 00	178 40 00	3,046	75	74		br. m.	do	
241	do	26 39 00	178 30 45	3,000	75	75	35.5	br. m.	do	
242	do	26 36 00	178 21 30	2,961	75	75		br. m.	do	
243	do	26 35 00	178 11 15	2,949	75	75	38.27	br. m.	do	

a "Marked "incorrect."

b "Incorrect."

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur-face.	Bot-tom.			
		° ' "	° ' "	Fath-oms.	°	°	°			
244	May 28	26 32 00	178 00 15	3,138	75	75	38?	br. m.	Red clay	
245	do	26 29 00	177 48 45	3,003	76	75		br. m.	do	
246	do	26 26 00	177 36 30	3,035	75	75	35	br. m.	do	
247	May 29	26 22 00	177 23 00	3,072	75	75		br. m.	do	
248	do	26 21 00	177 19 45	3,168	74	75		br. m.	do	
249	do	26 18 00	177 06 30	3,148	77	75	35	br. m.	do	
250	do	26 14 00	176 53 15	3,188	75	75		br. m.	do	
251	do	26 10 00	176 41 15	3,240	75	75	38.9?	br. m.	do	
252	do	26 06 00	176 29 30	3,230	75	76	35	br. m.	do	
253	do	26 01 00	176 16 00	3,252	77	76		br. m.	do	
254	do	25 57 00	176 04 45	3,242	75	75		br. m.	do	
255	do	25 57 00	175 53 00	3,240	75	75	35	br. m.	do	
256	May 30	25 50 00	175 42 30	3,260	75	75		br. m.	do	
257	do	25 47 00	175 32 00	3,250	75	76	60?	br. m. r. g.	do	
258	do	25 43 00	175 21 30	3,246	78	77	35.1	br. m. r. g.	do	
259	do	25 42 00	175 17 30	3,244	78	78		br. c.	do	
260	do	25 41 00	175 15 15	3,254	78	78		br. c.	do	
261	do	25 40 00	175 12 45	3,261	76	78	35	br. c.	do	
262	do	25 37 00	175 02 00	3,276	77	78		br. m.	do	
263	do	25 34 00	174 51 00	3,259	78	78		br. m.	do	
264	do	25 31 00	174 39 40	3,231	77	77	35	br. m.	do	
265	do	25 28 00	174 30 00	3,199	77	77		br. m.	do	
266	May 31	25 26 00	174 21 30	3,245	76	77	35	br. m.	do	
267	do	25 25 00	174 19 00	3,269	77	77		br. m.	do	
268	do	25 23 00	174 10 00	3,269	77	77		br. m.	do	
269	do	25 22 00	174 08 00	3,284	79	77		br. m.	do	
270	do	25 19 00	173 58 30	3,273	82	77	35	br. m.	do	
271	do	25 17 00	173 49 45	3,240	86	79		br. m.	do	
272	do	25 14 00	173 40 00	3,221	79	78		br. m.	do	
273	do	25 11 00	173 30 15	3,258	78	78		br. m.	do	
274	do	25 08 00	173 20 15	3,199	78	77	35	br. m.	do	
275	do	25 05 00	173 10 30	3,209	78	77		br. m.	do	
276	June 1	25 02 00	173 00 45	3,225	78	78	35	br. m.	do	
277	do	24 59 00	172 50 45	3,232	77	78		br. m.	do	
278	do	24 56 00	172 40 30	3,250	78	77	35	br. m.	do	
279	do	24 53 00	172 30 45	3,230	78	78		br. m.	do	
280	do	24 50 00	172 21 30	3,199	78	78	35	br. m.	do	
281	do	24 47 00	172 12 45	3,230	78	78		br. m.	do	
282	do	24 43 00	172 02 45	3,199	78	78		br. m.	do	
283	do	24 39 00	171 52 30	3,240	76	78	35	br. m.	do	
284	do	24 36 00	171 42 45	3,245	76	77		br. m.	do	
285	June 2	24 32 00	171 33 15	3,089	75	78	35	br. m. and r.	do	Manganese concretions.
286	do	24 31 00	171 31 15	3,187	76	78		br. m.	do	
287	do	24 27 00	171 21 15	3,250	75	77		br. m.	do	
288	do	24 22 00	171 12 00	3,334	74	78		br. m.	do	
289	do	24 18 00	171 03 00	3,339	73	78		br. m.	Red clay	
290	do	24 17 00	170 59 45	3,247	77	78		br. m.	do	
291	do	24 12 00	170 50 15	3,275	74	77		br. m.	do	
292	do	24 06 00	170 39 45	3,253	74	78		br. m.	do	
293	do	24 01 00	170 28 45	3,296	76	77		br. m.	do	
294	June 3	23 45 00	170 18 45	3,313	73	76		br. m.	do	
295	do	23 49 00	170 07 00	3,274	74	76		br. m.	do	
296	do	23 44 00	169 56 30	3,291	73	78		br. m.	do	
297	do	23 39 00	169 46 15	3,382	75	78		br. m.	do	
298	do	23 34 00	169 36 30	3,313	75	78		br. m.	do	
299	do	23 29 00	169 26 45	3,254	75	78		br. m.	do	
300	do	23 25 00	169 16 45	3,272	73	79		br. m.	Red clay	
301	do	23 20 00	169 07 15	3,242	76	75		br. m.	do	
302	June 4	23 16 00	168 57 30	3,207	75	77		br. m.	do	
303	do	23 11 00	168 48 00	3,207	76	79		br. m.	do	
304	do	23 07 00	168 38 15	3,176	76	79		br. m.	do	
305	do	23 02 00	168 28 30	3,214	79	79		lost	do	
306	do	22 57 00	168 18 15	3,285	78	78		br. m.	do	
307	do	22 54 00	168 08 30	3,275	76	78		br. m.	do	
308	do	22 51 00	167 57 50	3,381	77	78		lost	do	
309	do	22 48 00	167 47 15	2,992	76	78		lost	Red clay	
310	June 5	22 44 00	167 36 30	3,217	76	78		br. m.	do	
311	do	22 41 00	167 25 30	3,248	78	78		br. m.	do	
312	do	22 37 00	167 14 15	3,196	80	79	38.4	br. m.	do	
313	do	22 34 00	167 03 30	3,199	85	79		br. m.	do	
314	do	22 30 00	166 51 30	3,237	87	79	35	br. m.	do	Fragments of coscinodiscus rex.
315	do	22 27 00	166 40 00	3,261	81	79		br. m.	do	
316	do	22 25 00	166 28 30	3,261	78	79	35	br. m.	do	

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1899.	° ' "	° ' "	<i>Fathoms.</i>	°	°	°			
317	June 5	22 23 00	166 17 15	3,331	79	79		br. m.	Red clay...	
318	June 6	22 20 00	166 06 30	3,193	77	78	35	br. m.	do	
319	do	22 18 00	165 55 00	3,139	77	78		br. m.	do	
320	do	22 16 00	165 43 00	3,170	79	78	35	br. m.	do	
321	do	22 14 00	165 31 30	3,261	79	79		br. m.	do	
322	do	22 12 00	165 19 30	3,121	78	79	35	br. m.	do	
323	do	22 08 00	165 08 15	3,046	82	79		br. m.	do	
324	do	22 05 00	164 56 00	3,024	80	79	35	br. m.	do	
325	do	22 01 00	164 45 50	2,986	79	79		br. m.	do	
326	do	21 59 00	164 43 00	3,021	78	79		br. m.	do	
327	do	21 56 00	164 33 30	3,036	77	78	35	br. m.	do	
328	June 7	21 52 00	164 24 00	3,012	79	79		br. m. fn. spk	do	
329	do	21 48 00	164 15 00	2,993	77	79	35	br. m. fn. and	do	
								crs. sp.		
330	do	21 45 00	164 06 45	2,993	79	80		br. m. fn. and	do	
								crs. sp.		
331	do	21 44 00	164 04 00	2,997	80	81		br. m.	do	
332	do	21 39 00	163 54 45	2,988	77	80	35	br. m. blk. sp.	do	Phillipsite crystals.
333	do	21 35 00	163 45 15	2,965	78	80		br. m.	do	
334	do	21 32 00	163 35 30	2,902	76	80	35	br. m. fn. and	do	
								crs. sp.		
335	do	21 31 00	163 33 45	2,845	78	80		br. m. fn. and	do	
								crs. sp.		
336	do	21 27 00	163 23 45	2,424	79	80		br. m. r	do	Foraminifera.
337	do	21 26 00	163 21 30	2,287	79	80		br. m. r	do	
338	do	21 26 00	163 19 30	2,128	79	80	35	br. m. r	Globigerina ooze.	Large, manganese concretions.
										Do.
339	do	21 25 00	163 17 15	1,842	79	79		co. s.	do	Do.
340	do	21 24 00	163 15 00	1,447	76	78		crs. co. s. bk. sp.	do	Do.
341	do	21 24 00	163 14 15	1,315	76	78				No specimen.
342	do	21 24 00	163 13 15	1,380	77	78				Do.
343	June 8	21 23 00	163 11 15	1,298	77	78		co. s. bk. sp.	Globigerina ooze.	
344	do	21 23 00	163 10 15	1,228	77	78				Do.
345	do	21 22 00	163 09 10	1,173	78	78		crs. co. s. bk. sp.	Globigerina ooze.	
346	do	21 22 00	163 08 00	1,211	78	79		crs. co. s.	do	
347	do	21 21 00	163 07 00	1,215	78	79				Do.
348	do	21 21 00	163 04 45	1,606	78	79		crs. co. s.	Globigerina ooze.	
349	do	21 19 00	162 59 00	1,966	78	79		fn. co. s. br. m. fn. sp.	do	
350	do	21 15 00	162 48 30	2,240	80	81	35	br. m. co. s. r.	Red clay...	Fragments of Coscinodiscus rex.
										No specimen.
351	do	21 14 00	162 46 15	2,270	82	81				
352	do	21 12 00	162 39 45	2,568	87	81		br. m. crs. sp.	Red clay...	
353	do	21 10 00	162 34 45	2,825	77	81		br. m. crs. sp.	do	
354	do	21 06 00	162 23 45	2,836	82	81	35	br. m. crs. sp.	do	
								r.		
355	do	21 02 00	162 12 45	2,889	79	81		br. m. crs. sp.	do	
356	do	20 57 00	162 00 30	2,897	78	80	35	br. m. fn. sp.	do	
357	do	20 52 00	161 48 00	2,885	78	80		br. m. fn. sp.	do	
358	June 9	20 48 00	161 35 15	2,890	78	79	35	br. m. fn. sp.	do	
359	do	20 43 00	161 22 00	2,659	77	80		br. m. fn. sp.	do	
360	do	20 41 00	161 18 00	2,539	78	80		br. m. crs. sp.	do	
361	do	20 38 00	161 11 15	2,268	75	81		br. m. fn. co. s.	do	Few foraminifera.
362	do	20 36 00	161 05 15	1,937	76	81				Typical.
								co. s. and g.	Globigerina ooze.	
363	do	20 35 00	161 02 15	1,492	78	81	35, 3	crs. co. s. and g.	do	
364	do	20 32 00	161 02 00	1,723	81	81		crs. co. s. and g.	do	
365	do	20 28 00	161 01 30	1,601	82	81		crs. co. s.	do	
366	do	20 27 00	160 57 45	1,511	82	81		crs. co. s.	do	
367	do	20 26 00	160 54 00	1,251	81	80		Traces of r.	do	
368	do	20 26 00	160 51 45	1,013	82	82	36.4	Traces of r and s.	do	
369	do	20 25 00	160 49 45	966	81	81		crs. co. s.	do	
370	do	20 23 00	160 59 30	1,615	79	81				No specimen.
371	do	20 21 00	160 59 15	1,617	81	81				Do.

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
372	1899. June 9	20 18 00	160 58 45	<i>Fath- oms.</i> 1,738	80	81		fn. co. s.	Globigerina ooze.	
373	do	20 12 00	160 58 15	2,153	79	80		br. m. co. s. r.	Red clay	Manganese.
374	do	20 07 00	160 56 15	2,457	80	81		br. m. and r.	do	
375	do	20 03 00	160 52 45	2,509	80	80		br. m. and r.	do	
376	June 10	19 58 00	160 48 00	2,780	79	80		br. m. and r.	do	
377	do	19 52 00	160 38 45	2,611	79	80		br. m. and r.	do	
378	do	19 50 00	160 36 45	2,420	79	80		br. m. and r.	do	
379	do	19 44 00	160 33 15	2,203	80	80	35	br. m. and r.	do	No specimen.
380	do	19 38 00	160 32 15	2,124	78	80		br. m. and r.	do	Do.
381	do	19 38 00	160 30 15	1,846	78	80				
382	do	19 38 00	160 28 10	1,487	79	81		co. s.	Globigerina ooze.	
383	do	19 37 00	160 25 45	1,307	79	81		fn. co. s.	do	
384	do	19 37 00	160 23 15	747	79	81		co. s. and r.	do	Manganese.
385	do	19 37 00	160 21 45	720	79	81	37.9	crs. co. s.	do	
386	do	19 35 00	160 31 45	2,084	79	81		co. s. br. m. r.	do	
387	do	19 27 00	160 31 15	2,152	80	81	35	co. s. br. m. r.	do	
388	do	19 22 00	160 29 30	2,415	80	79		gy. br. m. r.	Red clay	
389	do	19 14 00	160 25 15	2,823	79	79	35	gy. br. m.	do	
390	do	19 10 00	160 21 45	3,006	80	79		gy. br. m. r.	do	
391	June 11	19 04 00	160 13 00	3,102	80	79	35	br. m.	do	
392	do	18 57 00	160 05 15	3,121	80	79		br. m.	do	
393	do	18 51 00	159 56 15	3,167	81	86	35	br. m.	do	
394	do	18 49 00	159 50 15	3,144	82	81		br. m.	do	
395	do	18 49 00	159 44 30	3,150	83	82		br. m.	do	
396	do	18 48 30	159 35 30	3,151	81	82	35	br. m.	do	
397	do	18 46 00	159 24 30	3,159	82	82		br. m.	do	
398	do	18 44 00	159 14 00	3,159	80	82	35.1	br. m.	do	
399	do	18 42 00	159 03 30	3,159	79	82		br. m.	do	
400	do	18 39 00	158 52 30	3,159	80	82	35	br. m.	do	
401	do	18 37 00	158 42 00	3,151	80	81		br. m.	do	
402	do	18 34 00	158 31 00	3,165	80	81	35	br. m.	do	
403	do	18 32 00	158 20 30	3,202	80	81		br. m.	do	
404	do	18 30 00	158 10 00	3,168	82	82	35	br. m.	do	
405	June 12	18 27 00	157 59 00	3,173	84	82		br. m.	do	
406	do	18 25 00	157 48 15	3,169	85	82	35	br. m.	do	
407	do	18 22 00	157 37 00	3,178	83	82		br. m.	do	
408	do	18 19 00	157 27 30	3,151	83	82		lost		No specimen.
409	do	18 17 00	157 18 00	3,129	81	82		br. m.	Red clay	
410	June 13	18 14 00	157 08 00	3,159	81	82		br. m.	do	
411	do	18 12 00	156 58 00	3,188	80	81	35	br. m.	do	
412	do	18 09 00	156 48 15	3,199	80	82		br. m.	do	
413	do	18 06 00	156 38 30	3,183	83	82		br. m.	do	
414	do	18 03 00	156 28 45	3,243	84	82		br. m.	do	
415	do	18 01 00	156 20 30	3,220	88	82	35	br. m.	do	
416	do	17 57 00	156 10 30	3,183	84	82		br. c.	do	
417	do	17 53 00	156 00 15	3,136	81	81	35	br. c.	do	
418	do	17 49 00	155 50 00	3,201	82	81		br. m.	do	
419	do	17 45 00	155 39 30	3,222	81	81	35	br. m. fn. sp.	do	
420	June 14	17 41 00	155 29 15	3,146	82	81		br. m. fn. sp.	do	
421	do	17 39 00	155 23 45	3,155	83	81		br. m.	do	
422	do	17 36 00	155 13 45	3,118	82	81	37.8	br. m.	do	
423	do	17 32 00	155 03 30	3,116	84	83		br. m.	do	
424	do	17 28 00	154 52 45	3,029	87	83	34.9	br. m. fn. sp.	do	
425	do	17 27 00	154 46 30	2,913	88	83		br. m. fn. sp.	do	
426	do	17 26 00	154 39 45	2,514	89	83		br. m. fn. sp.	do	
427	do	17 24 00	154 33 30	1,997	82	83		co. s. g.	Globigerina ooze.	
428	do	17 24 00	154 30 15	1,469	81	82	35.4			Do.
429	do	17 24 00	154 27 45	2,090	81	82				
430	do	17 24 00	154 38 15	2,292	80	81	34.9	r		A manganese nodule.
431	June 15	17 24 00	154 43 15	2,893	81	81		br. m. crs. sp.	Red clay	
432	do	17 18 00	154 43 15	2,921	79	81		br. m. fn. sp.	do	
433	do	17 12 00	154 42 30	2,947	79	81	35	br. m. fn. sp.	do	
434	do	17 08 00	154 40 00	2,796	82	82		br. c. fn. sp.	do	
435	do	17 05 00	154 36 15	2,638	83	82		br. c. crs. sp.	do	
436	do	17 02 00	154 32 00	2,346	84	83				No specimen.
437	do	16 59 00	154 28 30	1,593	83	83		fn. co. s.	Globigerina ooze.	
438	do	16 58 00	154 27 45	1,161	83	83				Do.
439	do	17 07 00	154 30 30	2,241	81	82	34.9	br. m. co. s.		Do.
440	do	17 15 00	154 34 45	2,598	86	82		br. m. fn. sp.	Red clay	
441	do	17 23 00	154 38 45	2,870	80	81		br. m. fn. sp.	do	

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
442	June 15	17 29 00	154 37 45	2,320	80	80	35	br. m. fn. co. s	Globigerina ooze.	
443	June 16	17 34 00	154 36 45	2,982	80	80		br. m. fn. sp.	Red clay	
444	do	17 39 00	154 34 30	3,047	80	80	34.9	br. m. fn. sp.	do	
445	do	17 43 00	154 29 45	3,041	80	80		br. m. fn. sp.	do	
446	do	17 45 00	154 24 15	3,036	81	82		br. m. fn. sp.	do	
447	do	17 45 00	154 17 30	3,058	84	82	36.3	br. c. fn. sp.	do	
448	do	17 43 00	154 05 15	3,110	84	83		br. m. fn. sp.	do	
449	do	17 42 00	154 01 15	3,129	87	83	36	br. m. fn. sp.	do	
450	do	17 39 00	153 50 15	3,154	86	83		br. m. fn. sp.	do	
451	do	17 35 00	153 39 30	3,158	84	82	35.8	br. m. fn. sp.	do	
452	do	17 32 00	153 28 45	3,067	82	82		br. c. fn. sp.	do	
453	do	17 29 00	153 17 45	2,363	81	82	35	br. m. fn. sp.	do	
454	do	17 34 00	153 16 45	2,375	81	82		br. m. crs. sp.	do	
455	June 17	17 38 00	153 16 15	2,446	80	80		br. m. fn. sp.	Red clay	No specimen.
456	do	17 29 00	153 19 45	2,466	79	80	35	br. m. crs. sp.	do	
457	do	17 24 00	153 18 45	2,353	80	81		co. s.	Globigerina ooze.	
458	do	17 22 00	153 13 45	1,466	82	82	35.8			
459	do	17 22 00	153 11 15	709	81	82		fn. co. s.	Globigerina ooze.	Do.
460	do	17 21 00	153 10 00	689	81	82				
461	do	17 21 00	153 08 45	711	81	82		crs. co. s.	do	
462	do	17 26 00	153 07 45	721	83	83	38.6	crs. co. s.	do	
463	do	17 31 00	153 06 45	1,913	80	82		co. s. and r.	do	
464	do	17 42 00	153 05 15	2,156	80	82	36			Do.
465	do	17 47 00	153 04 00	2,284	84	82		br. m. and fn. co. s.	Globigerina ooze.	
466	do	17 44 00	152 59 30	2,441	85	82		br. m. fn. sp.	do	
467	do	17 41 00	153 06 30	2,060	80	82		y. m. co. s. r.	do	Spherules of phillipsite in great number.
468	do	17 40 00	153 08 15	1,989	81	82		co. s.	do	
469	do	17 39 00	153 10 30	2,050	81	82	34.9	br. m. co. s.	do	
470	do	17 41 00	152 58 00	2,004	80	82		br. m. fn. sp.	Red clay	
471	do	17 39 00	152 51 45	2,325	81	82		br. m. co. s.	do	
472	June 18	17 36 00	152 45 45	2,000	80	81	35			No specimen.
473	do	17 33 00	152 40 00	1,679	79	81		co. s. bk. sp.	Globigerina ooze.	
474	do	17 33 00	152 35 45	1,159	79	81				Do.
475	do	17 33 00	152 34 15	1,309	79	79		crs. co. s.	Globigerina ooze.	
476	do	17 38 00	152 33 15	1,885	82	80				Do.
477	do	17 36 00	152 28 15	2,609	83	82	38.8	br. m. and r.	Red clay	
478	do	17 36 00	152 23 00	2,708	81	82		br. m. fn. sp.	do	
479	do	17 40 00	152 24 15	2,761	85	83	44.5	br. m. fn. and crs. sp.	do	
480	do	17 43 00	152 24 45	2,662	81	83		br. m. fn. sp.	do	
481	do	17 47 00	152 25 30	2,785	85	83		br. m. fn. sp.	do	
482	do	17 48 00	152 30 00	2,778	81	83	35.8	br. m. fn. sp.	do	
483	do	17 50 00	152 39 45	1,871	81	82		crs. co. s.	Globigerina ooze.	
484	do	17 51 00	152 44 30	2,371	81	82		br. m. fn. sp. and r.	Red clay	
485	do	17 51 00	152 55 30	2,741	80	81	35.2	br. m. fn. sp.	do	
486	June 19	17 51 00	153 06 30	2,506	81	82		br. m. fn. sp.	do	
487	do	17 45 00	153 16 15	2,615	80	81	35.9	br. m. fn. sp.	do	
488	do	17 39 00	153 26 30	3,015	82	83				Do.
489	do	17 34 00	153 36 15	3,148	82	83	40.2	br. m. fn. sp.	Red clay	
490	do	17 29 00	153 36 45	3,163	88	83		br. m. fn. sp.	do	
491	do	17 23 00	153 35 45	3,181	88	83	35.2			Do.
492	do	17 19 00	153 35 30	3,168	87	83				Do.
493	do	17 13 00	153 35 00	3,206	82	83				Do.
494	do	17 07 00	153 33 15	3,189	82	82		br. m. fn. sp.	Red clay	
495	do	16 57 00	153 29 30	3,204	83	81	35	br. m. crs. sp.	do	
496	June 20	16 52 00	153 27 50	3,204	83	82		br. m. crs. sp.	do	
497	do	16 44 00	153 22 00	3,190	82	82	35	br. m. fn. g.	do	
498	do	16 35 00	153 16 00	3,185	82	82		br. m. fn. g.	do	
499	do	16 28 00	153 08 00	3,192	81	82	35	br. m. fn. g.	do	
500	do	16 22 00	152 59 15	3,193	88	83		br. m. fn. sp.	do	
501	do	16 18 00	152 51 00	3,206	87	83	37.4	br. m. fn. sp.	do	
502	do	16 15 00	152 39 50	3,211	92	84		br. m. fn. sp.	do	
503	do	16 12 00	152 29 00	3,225	86	83		br. m. fn. sp.	do	

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
504	1899. June 20	16 09 00	152 18 15	Fath- oms. 3, 227	83	82	o			
505	do	16 06 00	152 07 45	2, 838	83	82	35	br. m. fn. sp.	Red clay	No specimen.
506	do	16 04 00	152 02 45	2, 169	83	82		r. fn. co. s. r.	Globigerina ooze.	
507	June 21	16 08 00	152 00 45	2, 372	82	82				Do.
508	do	16 11 00	151 58 45	2, 399	83	82		r		Do.
509	do	16 16 00	152 01 30	2, 680	81	83	34.9			Do.
510	do	15 55 00	152 03 00	2, 658	82	83				Do.
511	do	15 50 00	152 04 30	2, 003	83	83		crs. co. s		Do.
512	do	15 51 00	152 00 00	2, 368	81	83		br. m. r.	Red clay	Manganese.
513	do	15 53 00	151 55 30	2, 702	84	83	34.8	br. m. fn. sp.	do	
514	do	15 53 00	151 52 15	2, 893	87	83		br. m. fn. sp.	do	
515	do	15 50 00	151 45 45	3, 030	89	84				No specimen.
516	do	15 47 00	151 35 30	3, 252	85	84	35	br. m. fn. sp.	Red clay	
517	do	15 44 00	151 26 45	3, 227	84	83		br. m. fn. sp.	do	
518	do	15 41 00	151 16 45	3, 373	81	82	35	br. m. fn. sp.	do	
519	June 22	15 38 00	151 07 30	3, 273	83	82		br. m. fn. sp.	do	
520	do	15 35 00	150 58 30	3, 262	82	82	35	br. m. fn. sp.	do	
521	do	15 31 00	150 48 30	3, 254	82	82		br. m. fn. sp.	do	
522	do	15 28 00	150 38 30	3, 284	85	83	35	br. m. fn. sp.	do	
523	do	15 25 00	150 28 30	3, 201	88	83		br. m. crs. sp.	do	
524	do	15 22 00	150 19 30	3, 204	88	84	35	br. m. crs. sp.	do	
525	do	15 18 00	150 10 00	3, 211	88	83		br. m. crs. sp.	do	
526	do	15 14 00	149 58 45	3, 190	84	83	35	br. m. and g.	do	
527	do	15 10 00	149 48 30	3, 231	84	83				Do.
528	do	15 06 00	149 37 15	3, 120	83	83	35	br. m. and g.	Red clay	Do.
529	June 23	15 02 00	149 28 00	3, 175	82	83		br. m. and g.	do	
530	do	14 57 00	149 17 00	3, 118	83	82	35	br. m. and g.	do	
531	do	14 53 00	149 07 15	3, 105	82	83				Do.
532	do	14 48 00	148 57 15	3, 087	85	83	35			Do.
533	do	14 44 00	148 48 45	3, 087	90	84		br. m. and g.	Red clay	
534	do	14 41 00	148 41 00	3, 139	84	84	35	br. m. and g.	do	
535	do	14 36 00	148 30 00	3, 056	82	84		br. m. and g.	do	
536	do	14 35 00	148 26 15	2, 963	84	84		br. m. and g.	do	
537	do	14 33 00	148 20 15	3, 154	84	83	35	br. m. and r	do	
538	do	14 29 00	148 10 30	3, 098	82	83		br. m. and r	do	
539	do	14 28 00	148 07 45	2, 774	81	83				Do.
540	June 24	14 24 00	147 57 30	1, 888	80	82	35			Do.
541	do	14 23 00	147 55 15	1, 846	81	83		fn. co. s	Globigerina ooze.	
542	do	14 25 00	147 59 00	1, 996	81	82		fn. co. s	do	
543	do	14 29 00	147 59 45	1, 870	81	82				Do.
544	do	14 33 00	148 00 15	1, 946	80	83	34.9	crs. co. s. and g.	Globigerina ooze.	
545	do	14 21 00	148 03 00	1, 996	81	83				Do.
546	do	14 16 00	148 03 45	2, 414	80	84	35	crs. sp. s. and g.		Do.
547	do	14 13 00	148 03 45	2, 689	87	84		y. m. bk. sh.	Volcanic mud.	
548	do	14 15 00	148 06 45	3, 183	85	84		br. m. crs. sh. s.	Red clay	
549	do	14 10 00	147 58 15	1, 982	87	84				Do.
550	do	14 08 00	148 03 15	2, 975	82	84	35	br. m. and st.	Red clay	
551	do	14 04 00	148 01 45	3, 017	84	84		br. m. and st.	do	
552	do	14 00 00	147 59 15	2, 930	81	84		br. m. fn. sp.	do	
553	do	13 56 00	147 57 15	2, 712	81	84		br. m. fn. sp.	do	
554	do	13 52 00	147 56 00	3, 128	81	84		br. m. fn. sp.	do	
555	June 25	13 50 00	147 53 45	3, 154	84	82		br. m. fn. sp.	do	
556	do	13 47 00	147 49 30	3, 211	83	82	35	br. m. bk.	do	
557	do	13 46 00	147 41 00	3, 267	83	82		sh. s.		
558	do	13 46 00	147 33 30	3, 457	83	83	35.1	br. m. bk.		Do.
559	do	13 45 00	147 22 45	3, 658	90	84		br. m.	Diatom ooze.	
560	do	13 45 00	147 12 15	2, 558	88	85		co. s. and r.	Red clay	
561	do	13 53 00	147 18 45	3, 843	85	85	35.4			Do.
562	do	13 42 00	147 25 46	3, 945	83	85		br. m. sh. s.	Red clay	
563	do	13 38 00	147 24 00	3, 506	82	85	34.1			Do.
564	do	13 33 00	147 23 00	3, 461	83	85		br. m. sh. s.	Red clay	
565	do	13 28 00	147 22 15	3, 423	83	84		br. m. crs.	do	
566	June 26	13 23 00	147 20 00	3, 389	82	83		sh. s.		
567	do	13 18 00	147 16 45	3, 379	82	83		br. m. crs.	do	

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
568	June 26	13 14 00	147 13 45	3,379	83	82	br. m. crs. sp.	Red clay...	
569do.....	13 14 00	147 08 15	3,190	86	83	35	br. m. crs. sh. s.do.....	
570do.....	13 14 00	147 02 15	3,057	86	84	br. m. and p.do.....	
571do.....	13 14 00	146 56 30	3,288	89	85do.....	No specimen.
572do.....	13 13 30	146 50 45	4,085	90	85	35.4	Red clay...	
573do.....	13 24 00	146 49 45	4,547	86	85do.....	Do.
574do.....	13 34 00	146 50 45	4,913	82	83do.....	Do.
575	June 27	13 45 00	146 51 30	4,563	82	83	br. m. crs. sp.	Red clay...	
576do.....	13 55 00	146 52 30	4,490	82	84	br. m. and st.do.....	
577do.....	14 05 00	146 53 45	3,897	82	85	gy. m. fn. s.do.....	
578do.....	14 08 00	147 03 45	4,563	79	85	gy. m.do.....	
579do.....	14 11 00	147 14 45	4,618	82	85do.....	Do.
580do.....	14 14 00	147 26 45	3,895	82	83	y. m. crs. sp.	Red clay...	
581	June 28	14 18 00	147 38 45	1,848	82	83do.....	Do.
582do.....	14 19 00	147 40 45	1,686	82	84do.....	Do.
583do.....	14 19 00	147 43 15	1,631	83	84	co. s.	Globigerina ooze.	
584do.....	14 29 00	147 42 30	1,945	82	84do.....	Do.
585do.....	14 39 00	147 41 45	2,604	83	84	34.9	br. m. crs. sp.	Red clay...	
586do.....	14 49 00	147 41 15	3,683	80	83do.....	Do.
587do.....	14 49 30	147 42 00	3,534	82	84do.....	Do.
588do.....	14 56 00	147 48 10	3,263	83	84	br. m. crs. s.	Red clay...	
589do.....	15 06 00	147 46 20	3,150	82	85do.....	Do.
590do.....	15 15 00	147 40 20	3,607	82	84do.....	Do.
591	June 29	15 25 00	147 38 30	4,204	78	82	br. m. crs. sp.do.....	
592do.....	15 35 00	147 36 45	3,832	79	84	br. m. fn. sp.do.....	
593do.....	15 44 30	147 35 00	3,404	83	85	hrd. c. fn. s.	Volcanic mud.	
594do.....	15 44 00	147 24 30	2,233	85	85	br. m. bk. s.do.....	
595do.....	15 44 00	147 18 45	2,409	89	85	br. m. bk. s.do.....	
596do.....	15 44 00	147 09 00	2,124	88	85	dk. gy. s.do.....	
597do.....	15 40 00	146 59 30	2,941	86	85	34.9	bk. sandbr.do.....	
598do.....	15 37 00	146 49 45	3,130	83	85	m.do.....	
599do.....	15 33 00	146 40 30	2,976	84	85	br. m.do.....	Do.
600do.....	15 30 00	146 31 15	2,536	83	84	br. m. co. s.	Volcanic mud.	
601	June 30	15 26 00	146 21 45	2,178	79	84	35	fn. dk. s. sh.do.....	
602do.....	15 22 00	146 12 30	1,771	81	84do.....	Do.
603do.....	15 15 00	146 07 30	1,743	81	83	bk. and gy.	Volcanic mud.	
604do.....	15 49 00	147 32 30	2,951	83	84	br. m. and fn. bk. s.do.....	
605do.....	15 52 00	147 40 30	3,378	83	85	grn. c. and s.do.....	
606do.....	15 55 00	147 48 30	3,360	83	84do.....	Do.
607	July 1	15 57 00	147 56 00	2,846	83	84do.....	Do.
608do.....	15 57 00	148 04 00	2,969	82	84do.....	Do.
609do.....	15 54 00	148 11 15	2,841	83	85	35	br. m. bk. s.	Red clay...	
610do.....	15 47 00	148 18 08	1,780	87	85do.....	Do.
611do.....	15 41 00	148 22 15	2,409	83	85	34.9	br. m. bk. s.do.....	Do.
612do.....	15 33 30	148 27 00	2,369	82	85	bk. and wh. s.	Volcanic sand.	
613do.....	15 26 00	148 31 15	1,092	82	85	36.1	red and wh. s.	Globigerina ooze.	
614do.....	15 19 30	148 39 00	3,230	82	84	br. c. bk. s.	Red clay...	
615do.....	15 11 00	148 38 00	3,178	80	83	br. m.do.....	
616do.....	15 01 30	148 37 00	3,077	78	82	br. m. sh.do.....	
617	July 2	14 53 00	148 36 00	2,987	80	83	35	br. m. sh.do.....	
618do.....	15 17 00	148 30 50	2,022	80	84	wh. s. bk. sp.	Globigerina ooze.	
619do.....	15 15 00	148 27 45	2,414	81	84do.....	Do.
620do.....	15 11 00	148 27 20	2,567	88	84	wh. s. bk. sp.do.....	Do.
621do.....	15 13 00	148 23 00	2,555	88	84	br. m. and s.	Red clay...	
622do.....	15 14 30	148 19 00	2,537	88	85	br. m. and bk. s.do.....	
623do.....	15 19 00	148 15 00	2,088	85	86	br. m. and bk. s.	Globigerina ooze.	
624do.....	15 19 00	148 10 30	2,414	87	85	35	br. m. and bk. s.	Red clay...	
625do.....	15 20 30	148 06 08	2,578	86	85	bk. s.do.....	
626do.....	15 24 00	148 01 30	2,968	77	84	35	bk. s. fn. g.do.....	Do.
627do.....	15 29 00	147 58 45	3,158	81	84do.....	Do.
628do.....	15 38 00	147 54 00	3,381	83	84	35.1	br. m.	Red clay...	

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1899.	° ' "	° ' "	Fathoms.	°	°	°			
629	July 3	15 46 00	147 49 30	3,302	83	84		br. m.	Red clay...	
630	do	15 46 00	147 29 15	2,339	82	83	35	br. m.	do	
631	do	15 46 00	147 14 30	2,253	81	84		fn. co. s.	Globigerina ooze.	
632	do	15 43 00	147 04 50	2,559	83	84	35.3	br. m. bk. s.	Volcanic mud.	
633	do	15 19 00	146 16 30	2,052	87	85	35	br. m. bk. s.	do	
634	do	15 10 00	146 15 15	2,154	83	85		bk. s. co. s.	do	
635	do	15 00 00	146 14 15	2,285	83	85	35	bk. s. co. s.	do	
636	do	14 51 00	146 11 08	2,360	82	84		br. m. bk. s.	do	
637	do	14 41 00	146 06 00	2,352	82	84	35	bk. s. co. s.	do	
638	July 4	14 32 00	146 02 00	2,342	81	84		bk. s. br. m.	do	
639	do	14 22 00	145 57 15	2,297	80	84	35	bk. s. br. m.	do	
640	do	14 12 00	145 52 40	2,238	81	83		br. m. fn.	do	
								bk. s.		
641	do	14 04 00	145 48 00	2,187	80	83	35	bk. s. co. s.	do	
642	do	13 54 00	145 43 30	2,014	82	84		br. m. bk. s.	Globigerina ooze.	
643	do	13 45 00	145 38 45	1,757	81	85	35	co. s. bk. s.	do	No specimen.
644	do	13 39 00	145 29 00	1,483	82	85				
645	do	13 38 00	145 16 00	1,102	83	86	36	co. s.	Globigerina ooze.	
646	do	13 37 00	145 05 00	648	80	86		co. s.	do	
647	do	13 32 00	145 02 00	605	80	85	39.6	co. s.	do	
648	do	13 26 00	144 59 00	720	80	85		fn. co. s.	do	
649	do	13 21 00	144 57 00	762	80	85		fn. co. s.	do	
650	do	13 16 00	144 54 00	768	80	84		fn. co. s.	do	
651	do	13 10 00	144 52 00	907	80	85	37	co. s. bk. s.	do	
652	do	13 05 00	144 50 00	998	79	84		co. s.	do	
653	July 5	13 17 30	145 07 00	1,137	80	84	36	fn. co. s.	do	
654	do	13 18 30	144 53 00	579	81	84				Do.
655	do	13 18 00	144 51 00	480	81	84				Do.
656	do	13 18 00	144 49 00	404	82	85				Do.
657	do	13 18 00	144 48 00	304	82	85				Do.
658	do	13 18 00	144 47 30	208	82	85				Do.
659	do	13 18 00	144 47 15	85	82	85				Do.
660	do	13 05 00	144 41 00	709	84	85				Do.
661	do	13 11 30	144 32 00	812	85	88	37.2	co. s.	Globigerina ooze.	
662	do	13 13 25	144 32 30	1,260	83	88				

GUAM TO LUZON.

663	July 7	13 26 30	144 36 30	457	83	80		co. s. sh. bk. s.	Coral sand.	No specimen.
664	do	13 27 00	144 35 00	1,016	83	85		co. s. wh. s.	Volcanic mud.	
665	do	13 27 00	144 25 45	1,652	82	84	35.2	yl. m. co. s.	do	
666	do	13 27 00	144 23 30	1,693	81	84		co. s. bk. s.	do	
667	do	13 27 30	144 13 15	2,009	79	84				Do.
668	July 8	13 27 30	144 10 30	2,094	81	84				
669	do	13 28 00	144 00 00	1,696	82	83	35.2			Do.
670	do	13 28 00	143 57 15	1,376	82	83		bk. s.	Volcanic mud.	
671	do	13 28 00	143 54 45	1,415	82	83		co. s. bk. s.	do	Do.
672	do	13 28 00	143 52 30	1,820	83	83		br. m. bk. and co. s.	do	
673	do	13 28 20	143 42 45	1,967	82	83	35	lt. br. m. bk. s.	do	
674	do	13 28 30	143 40 20	1,862	82	85		yl. bro. m. bk. s.	do	
675	do	13 29 00	143 29 00	2,007	86	85		yl. bro. m. bk. s.	do	Do.
676	do	13 29 00	143 27 15	1,811	88	86				
677	do	13 29 00	143 22 15	1,946	84	86	35.2	yl. br. m. fn. bk. sp.	Volcanic mud.	
678	do	13 30 30	143 11 30	1,883	84	86		br. m. fn. bk. sp.	do	
679	do	13 31 00	143 09 00	2,310	84	85		br. m. fn. bk. sp.	do	Fine glass; much manganese.
680	do	13 32 30	142 57 30	2,010	84	85		br. m. lava ..	do	

^a Soundings 646 to 662 taken in vicinity of Port Tarafoto, thence to San Luis d'Apra Harbor, Island of Guam.

Abstract of the official record of soundings—Continued.

GUAM TO LUZON—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
681	July 8	13 33 00	142 54 30	2,648	82	84	35.6			No specimen.
682	do	13 36 30	142 55 30	2,650	82	84				Do.
683	do	13 28 30	142 55 30	2,319	83	84		br. m. fn. bk. sp.	Red clay	
684	do	13 28 30	142 52 15	2,514	83	84		br. m.	do	
685	July 9	13 30 00	142 40 45	2,009	83	85	35.4			Do.
686	do	13 30 30	142 37 20	1,586	82	84				Do.
687	do	13 31 00	142 33 45	1,553	82	85				Do.
688	do	13 31 30	142 30 15	1,346	83	85		bk. and wh. s. lava.	Globigerina ooze.	Manganese.
689	do	13 32 00	142 26 45	1,569	82	85		wh. s. bk. sp.	do	
690	do	13 33 30	142 15 15	1,863	84	85	35.3			No specimen.
691	do	13 34 00	142 12 00	1,841	83	85		yl. m. and s.	Globigerina ooze.	
692	do	13 35 30	142 00 30	1,739	87	86				Do.
693	do	13 36 00	141 57 30	1,977	85	86				Do.
694	do	13 37 30	141 47 00	2,392	88	86				Do.
695	do	13 38 00	141 44 15	2,349	89	86	36			Do.
696	do	13 40 00	141 34 00	2,514	86	86				Do.
697	do	13 40 30	141 32 15	2,506	80	85		br. m.	Red clay	
698	do	13 43 00	141 20 45	2,591	82	85	35.2	yl. m. s.	do	
699	do	13 43 00	141 18 15	2,632	81	85				Do.
700	July 10	13 46 00	141 08 00	2,663	80	84				Do.
701	do	13 46 00	141 05 15	2,691	82	85				Do.
702	do	13 49 00	140 55 15	2,691	81	85				Do.
703	do	13 49 30	140 52 45	2,710	82	85				Do.
704	do	13 52 00	140 43 15	2,740	78	84	35.1	br. m.	Red clay	
705	do	13 54 30	140 33 15	2,710	78	85		br. m.	do	
706	do	13 56 00	140 23 15	2,726	75	85	35			Do.
707	do	13 58 00	140 13 15	2,726	77	84		br. m.	Red clay	
708	do	14 00 00	140 03 15	2,647	79	84	35	br. m.	do	
709	do	14 01 30	139 52 45	2,375	76	84		br. m. fn. sp.	do	
710	do	14 02 00	139 50 30	2,231	77	84				Do.
711	do	14 03 00	139 45 20	2,317	77	84	35	br. m. ers. sp.	Red clay	Do.
712	July 11	14 04 00	139 40 45	2,099	77	84		yl. bro. m.	do	
713	do	14 04 00	139 38 15	2,212	81	84				Do.
714	do	14 06 00	139 28 30	2,992	81	84	35			Do.
715	do	14 08 00	139 17 45	2,639	82	84.5		br. m. fn. and ers. sp.	Red clay	
716	do	14 08 00	139 14 45	2,596	80	85		br. m. fn. bk. sp.	do	
717	do	14 10 00	139 03 45	2,837	81	85		br. m. fn. bk. sp.	do	
718	do	14 10 00	139 02 00	2,674	83	85		br. m. fn. bk. sp.	do	
719	do	14 12 00	138 54 45	2,705	82	85	67.6	hd. c.		Do.
720	do	14 13 30	138 47 30	2,374	81	85				Do.
721	do	14 14 00	138 45 45	2,519	81	85		br. m. fn. sp.	Red clay	
722	do	14 15 30	138 38 45	2,476	81	84	67	br. m.	do	
723	July 12	14 17 00	138 31 30	2,605	80	84	67			Do.
724	do	14 17 30	138 29 30	2,839	82	84		br. m. hd. c.	Red clay	
725	do	14 19 00	138 23 15	2,596	82	84		br. m. hd. c.	do	
726	do	14 19 30	138 21 00	2,638	82	84		br. m. hd. c.	do	
727	do	14 21 00	138 13 30	2,959	84	85				Do.
728	do	14 23 00	138 04 00	2,797	84	85				Do.
729	do	14 24 00	137 55 00	2,704	83	85		br. m.	Red clay	
730	do	14 25 30	137 45 45	2,761	86	84		br. m.	do	
731	do	14 27 00	137 36 30	2,782	82	84		br. m.	do	
732	do	14 28 00	137 27 45	2,568	79	84		br. m.	do	
733	July 13	14 28 30	137 25 00	2,477	79	84				Do.
734	do	14 29 00	137 21 10	2,477	82	84		br. m. bk. sp.	Red clay	
735	do	14 30 00	137 15 30	2,677	80	84		br. m. bk. sp.	do	
736	do	14 31 00	137 05 45	2,602	82	84.5		br. m. bk. sp.	do	
737	do	14 33 00	136 56 45	2,652	79	85		br. m.	do	
738	do	14 34 00	136 48 00	2,870	83	85		br. and yl. m. and c.	do	
739	do	14 33 00	136 40 10	2,862	83	84		br. m.	do	
740	do	14 32 00	136 30 15	2,735	82	84		br. c.	do	
741	do	14 30 30	136 20 00	2,907	81	83				Do.
742	do	14 29 00	136 10 30	3,145	77	83		gy. gu. m.	Red clay	
743	July 14	14 28 00	136 00 15	3,118	77	83		bl. gu. m.	Diatom ooze.	
744	do	14 26 30	135 50 30	2,879	82	83		gr. gu. m.	do	
745	do	14 25 00	135 40 30	2,617	81	84		br. m.	Red clay	
746	do	14 24 00	135 31 00	2,788	83	84		br. gu. m.	Diatom ooze.	

Abstract of the official record of soundings—Continued.

GUAM TO LUZON—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
747	July 14	14 23 00	135 21 00	2,731	85	85	br. m.	Red clay...	
748do....	14 24 00	135 10 10	2,891	82	84	br. m.do....	
749do....	14 24 30	135 00 45	2,819	82	84	br. m.	Diatom ooze.	
750do....	14 25 00	134 51 30	2,679	82	83	gy. m.do....	
751	July 15	14 25 30	134 42 30	2,679	82	83	br. m.	Red clay...	
752do....	14 26 00	134 34 00	2,432	83	83	bl. gn. m.	Diatom ooze.	
753do....	14 26 00	134 31 30	1,913	80	84	br. s.	Globigerina ooze.	
754do....	14 26 00	134 29 15	1,937	80	84	br. c. and s.do....	
755do....	14 26 00	134 27 00	1,935	80	84.5	wh. s. br. m.do....	
756	July 24	14 26 00	134 30 45	2,307	81	82	br. m. and s.do....	
757do....	14 26 15	134 26 30	2,158	81	82	yl. m. and s.do....	
758	July 25	14 26 30	134 23 00	1,780	81	84	yl. m. and s.do....	
759do....	14 26 45	134 20 00	1,657	81	84	yl. m. and s.do....	
760do....	14 27 00	134 17 00	1,560	81	84	yl. m. and s.do....	
761do....	14 27 15	134 13 45	1,619	81	84	sh. s. lavado....	
762do....	14 27 40	134 11 00	1,782	80	84	35.2	fn. wh. s.do....	
763do....	14 28 00	134 05 30	2,072	85	85	br. m. fn. bk.	Red clay...	
764do....	14 29 00	133 56 15	2,487	82	85	br. m.	Diatom ooze.	
765do....	14 29 45	133 47 00	2,688	83	85	br. m.	Red clay...	
766do....	14 30 00	133 40 15	2,799	82	86	br. m.do....	
767do....	14 31 00	133 33 45	2,827	81	86	35.4	br. m.do....	Diatoms.
768do....	14 32 00	133 23 00	2,988	86	86do....	No specimen.
769do....	14 33 30	133 12 00	2,914	83	85do....	Do.
770do....	14 35 00	133 01 00	2,888	81	85	35.3	br. m.	Red clay...	Do.
771	July 26	14 36 00	132 50 00	2,951	80	84do....	Do.
772do....	14 38 00	132 39 00	3,344	82	83	35.8do....	
773do....	14 39 00	132 28 00	3,119	81	83	br. c.	Red clay...	
774do....	14 41 00	132 17 00	3,029	83	84	35.5	br. m.do....	
775do....	14 42 00	132 06 45	3,423	88	85	br. m.do....	
776do....	14 43 30	131 55 45	3,283	86	85	br. and gn.	Diatom ooze.	
777do....	14 45 00	131 45 30	3,421	84	85	lt. br. m.	Red clay...	
778do....	14 46 00	131 34 45	3,089	85	86	35.5	br. m.do....	
779do....	14 47 00	131 24 15	3,172	83	85	br. m.do....	
780do....	14 47 30	131 13 30	3,354	83	85	35.4	br. m.do....	
781do....	14 48 30	131 03 00	3,252	81	84	wh. and br.	Diatom ooze.	
782do....	14 49 00	130 52 30	3,129	81	84	35.3	br. m.	Red clay...	
783do....	14 50 00	130 42 00	3,264	82	83	br. m.do....	
784	July 27	14 50 00	130 31 30	3,547	83	84	35.7	gy. m.	Diatom ooze.	
785do....	14 50 30	130 20 45	3,237	87	85	br. m.	Red clay...	
786do....	14 51 00	130 09 45	3,148	88	85	35.6	br. m.do....	
787do....	14 52 00	129 57 00	3,175	90	86	br. m.do....	
788do....	14 53 30	129 45 15	3,318	84	86	35.6	br. m.do....	
789do....	14 55 00	129 34 15	3,041	82	85	br. m.do....	
790do....	14 56 30	129 23 15	3,119	82	85	32	br. m.do....	
791do....	14 58 00	129 12 15	3,011	82	84	br. m.do....	
792do....	15 00 00	129 02 00	3,158	80	84	35.5	br. m. and st.do....	
793do....	15 02 00	128 52 00	3,099	81	84	br. m.do....	
794do....	15 04 30	128 41 40	2,840	82	84	35.5	br. m.do....	
795do....	15 06 30	128 31 30	3,093	85	86	br. m.do....	
796do....	15 08 30	128 22 45	2,670	85	86	br. m.do....	
797do....	15 09 00	128 20 00	2,767	84	86	br. m.do....	
798do....	15 10 00	128 09 30	3,098	85	86	36	br. m.do....	
799do....	15 10 00	127 59 15	3,025	86	86	br. m.do....	
800	July 28	15 10 30	127 49 30	3,108	82	85	35.5	br. m. and g.do....	
801do....	15 10 00	127 40 45	3,298	82	85	br. m.do....	
802do....	15 09 30	127 31 40	2,844	82	84	br. m.do....	
803	July 29	15 09 00	127 22 30	2,943	82	84	35.3	br. m.do....	
804do....	15 08 30	127 13 20	2,995	81	84	br. m.do....	
805do....	15 08 00	127 04 15	3,026	82	85	35.4	br. m.do....	
806do....	15 07 00	126 54 45	2,929	83	85	br. m.do....	
807do....	15 06 00	126 44 45	3,121	86	86	br. m.do....	
808do....	15 05 00	126 36 30	2,855	85	86	br. m.do....	
809do....	15 05 30	126 27 00	3,134	84	86	22.9	br. m.do....	
810do....	15 06 00	126 17 45	3,252	84	86	br. m.do....	
811do....	15 06 00	126 08 00	3,047	83	86	36	br. m.do....	
812do....	15 06 00	125 58 00	3,130	83	86	br. m.do....	

Abstract of the official record of soundings—Continued.

GUAM TO LUZON—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1899.	° ' "	° ' "	<i>Fathoms.</i>	°	°	°			
813	July 29	15 06 30	125 48 15	2,819	83	86	35.2	br. m.	Red clay...	
814	July 30	15 07 00	125 38 15	3,144	83	84	br. m.	do	
815	do	15 07 00	125 28 30	2,792	82	84	lost	do	
816	do	15 08 00	125 18 45	2,936	84	85	br. m.	do	
817	do	15 08 00	125 08 45	2,911	82	85	br. m.	do	
818	do	15 08 00	124 58 45	3,182	88	87	lt. br. m.	do	
819	do	15 08 00	124 54 30	3,116	86	86	35.7	br. m.	do	
820	do	15 09 00	124 44 00	2,817	87	87	br. c.	do	
821	do	15 10 00	124 33 30	2,468	83	86	35.4	br. c.	do	
822	do	15 10 30	124 30 15	2,427	85	86	br. c.	do	
823	do	15 12 00	124 20 00	2,683	83	85	br. c. and m.	do	
824	do	15 14 00	124 09 20	2,994	83	86	br. c.	do	
825	July 31	15 15 00	123 58 45	2,771	82	85	lt. br. m. fn. s.	do	
826	do	15 17 00	123 48 15	2,360	83	85	35.4	br. m. fn. s.	do	
827	do	15 19 00	123 38 00	1,401	82	85	br. m. fn. s.	Blue mud...	
828	do	15 20 00	123 34 30	1,390	81	84	br. m. fn. s.	do	
829	do	15 21 00	123 28 45	1,411	82	84	br. m. fn. s.	do	
830	do	15 22 30	123 22 45	1,515	82	85	br. m. fn. s.	do	
831	do	15 25 00	123 10 50	2,100	83	86	35.3	br. m. fn. s.	do	
832	do	15 28 30	122 58 40	2,458	85	86	br. m. fn. s.	do	
833	do	15 30 00	122 51 30	2,740	83	86	35.6	lt. br. m.	do	
834	do	15 29 30	122 40 45	2,600	85	87	br. m. fn. bk.	do	
835	do	15 28 30	122 29 50	2,259	86	87	s.	do	
836	do	15 27 00	122 19 45	1,364	86	89	35.6	lt. br. m. fn.	do	
837	do	15 26 00	122 16 15	1,286	84	88	bk. sp.	do	
838	do	15 25 00	122 12 40	1,406	84	87	br. m. and s.	do	
839	do	15 24 00	122 10 15	1,395	83	87	br. m. and s.	do	
840	do	15 23 00	122 08 00	1,478	84	87	br. m. and s.	do	
841	do	15 22 30	122 05 45	1,498	83	87	gn. c.	do	
842	Aug. 1	15 22 00	122 03 30	1,330	83	86	br. m. and s.	Blue mud...	
843	do	15 21 00	122 01 15	1,449	88	86	gn. and br. m.	do	
844	do	15 20 00	121 59 15	1,449	89	86	35.8	gn. and br. m.	do	
845	do	15 19 00	121 56 20	1,459	86	86	br. m.	do	
846	do	15 18 00	121 54 00	1,463	80	86	gn. m.	Green mud...	
847	do	15 16 30	121 49 15	1,481	81	86	gn. m.	do	
848	do	15 16 30	121 44 45	1,101	84	86	gn. m.	do	
849	do	15 16 00	121 40 00	737	82	86	gn. m.	do	
850	do	15 17 00	121 34 45	157	83	86	gn. m.	do	
851	do	15 17 00	121 34 00	134	83	86	gn. m.	do	No specimen.
852	do	15 17 15	121 33 00	120	83	86	gn. m.	do	Do.
853	do	15 17 30	121 31 45	103	83	86	gn. m.	do	No specimen.

LUZON TO GUAM.

854	Aug. 19	15 09 00	121 37 30	180	81	83	gr. c.		
855	do	15 26 30	121 47 15	1,599	81	85	gr. br. m.	Green mud...	
856	do	15 12 00	121 57 30	829	83	85	gr. c.	do	
857	do	15 31 30	122 08 00	1,046	83	85	gr. c.	do	
858	do	15 17 00	122 18 15	1,458	82	84	gr. and br.	do	
859	do	15 35 00	122 29 00	2,390	79	84	m. bk. sps.	do	
860	Aug. 20	15 19 15	122 40 15	2,090	80	83	br. m.	do	
861	do	15 35 15	122 52 45	3,083	79	82	br. m.	do	No specimen.
862	do	15 16 00	123 03 00	1,550	79	85	br. m.	Green mud...	
863	do	15 30 00	123 17 15	2,424	78	85	br. m.	do	
864	do	15 12 20	123 25 00	821	81	85	br. m.	do	
865	do	15 26 30	123 40 30	2,058	82	84	br. m.	do	
866	do	15 09 45	123 50 30	1,272	81	84	br. m.	Blue mud...	Do.
867	Aug. 21	15 24 00	124 05 15	2,985	83	84	br. m.	Red clay...	
868	do	15 07 15	124 14 00	2,136	82	83	br. m.	Green mud...	
869	do	15 21 30	124 28 15	2,440	83	85	br. m.	Red clay...	
870	do	15 04 00	124 34 45	3,140	85	85	br. m.	do	
871	do	15 19 15	124 44 45	2,348	84	86	35.2	br. m.	do	
872	do	15 01 00	124 53 00	3,260	85	84	br. m.	do	
873	do	15 19 00	125 04 20	2,920	85	85	35.4	br. m.	do	
874	Aug. 22	15 01 30	125 13 00	2,988	83	83	br. m.	do	
875	do	15 19 15	125 25 15	2,573	85	84	35.4	br. m.	do	
876	do	15 01 30	125 36 30	2,541	87	86	br. m.	do	

Abstract of the official record of soundings—Continued.

LUZON TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
877	Aug. 22	15 14 30	125 44 30	2,862	89	86	35.4	br. m.	Red clay	
878	do	14 57 00	125 54 00	2,957	84	86	br. m.	do	
879	do	15 14 00	126 02 30	2,990	85	85	35.6	br. m.	do	
880	Aug. 23	14 55 45	126 10 00	2,961	81	84	br. m.	do	
881	do	15 13 30	126 19 15	3,312	82	83	br. m.	do	
882	do	14 55 15	126 27 40	2,946	79	84	35.6	br. m.	do	
883	do	15 12 45	126 39 00	2,711	81	84	br. m.	do	
884	do	14 57 40	126 47 30	3,310	82	85	35.7	br. m.	do	
885	do	15 15 45	126 58 40	2,909	81	83	br. m.	do	
886	do	14 58 15	127 09 15	3,048	82	83	35.6	br. m.	do	
887	Aug. 24	15 17 00	127 19 30	2,943	81	84	br. m.	do	
888	do	14 59 30	127 31 00	3,009	80	84	35.5	br. m.	do	
889	do	15 17 00	127 40 45	3,006	81	84	br. c.	do	
890	do	15 00 00	127 54 00	3,014	86	85	35.5	br. m.	do	
891	do	15 17 45	128 01 30	2,987	90	85	br. m.	do	
892	do	14 59 15	128 11 45	3,234	83	84	35.7	br. m.	do	
893	do	15 16 00	128 18 15	3,145	82	84	br. m.	do	
894	Aug. 25	14 57 20	128 22 15	2,889	80	83	br. m.	do	
895	do	15 14 00	128 30 30	3,030	82	84	35.6	br. m.	do	
896	do	14 52 00	128 37 30	3,342	br. m.	do	
897	do	15 12 30	128 50 30	3,189	82	85	35.6	br. m.	do	
898	do	14 53 15	129 01 15	3,346	82	84	br. m.	do	
899	Aug. 26	15 10 00	129 16 30	2,864	81	83	35.5	br. m.	do	
900	do	14 53 00	129 27 40	3,159	br. m.	do	
901	do	15 11 15	129 37 30	2,781	87	85	35.6	br. m.	do	
902	do	14 45 40	129 50 00	2,945	88	80	br. m.	do	
903	do	15 00 00	129 55 30	2,809	83	85	35.5	br. m.	do	
904	do	14 39 30	130 05 30	3,096	83	83	br. m.	do	No specimen.
905	Aug. 27	14 55 00	130 13 00	3,128	80	83	br. m.	Red clay	
906	do	14 36 00	130 23 00	3,204	80	84	br. m.	do	
907	do	14 51 30	130 32 30	3,490	86	84	br. m.	do	
908	do	14 37 30	130 41 30	3,132	88	85	35.8	br. m.	do	
909	do	14 54 45	130 15 15	3,125	88	86	br. m.	do	
910	do	14 38 00	131 03 00	2,969	82	85	35.8	br. m.	do	
911	do	14 56 00	131 14 45	3,295	82	84	br. m.	do	
912	Aug. 28	14 40 00	131 26 40	2,985	80	83	35.9	br. m.	do	
913	do	14 57 30	131 39 00	2,823	81	84	br. m.	do	
914	do	14 42 15	131 51 45	3,065	83	84	35.5	br. m.	do	
915	do	14 55 00	132 00 30	3,118	87	85	br. m.	do	
916	do	14 35 30	132 09 30	3,103	85	85	35.9	br. m.	do	
917	do	14 49 45	132 20 00	3,246	81	85	br. m.	do	
918	do	14 32 00	132 26 00	3,253	80	84	35.8	br. m.	do	
919	Aug. 29	14 48 45	132 37 20	2,998	79	83	br. m.	do	
920	do	14 31 15	132 42 30	3,327	80	84	35.8	br. and gy. m.	Diatom ooze	
921	do	14 49 15	132 54 00	2,499	86	85	br. m.	Red clay	
922	do	14 32 00	133 00 30	2,769	88	85	35.6	br. m.	do	
923	do	14 44 00	133 11 45	2,322	85	86	br. m.	do	
924	do	14 23 30	133 20 15	2,878	81	85	35.8	br. m.	do	
925	do	14 39 30	133 32 30	2,905	83	84	br. m.	do	
926	Aug. 30	14 20 30	133 41 00	2,851	78	83	35.6	br. m.	do	
927	do	14 36 30	133 54 00	2,494	79	83	br. m.	do	
928	do	14 17 00	134 01 45	2,593	83	84	35.7	br. m.	do	
929	do	14 36 00	134 16 07	1,964	83	85	gy. m. and s.	Globigerina ooze.	
930	do	14 17 15	134 23 45	1,850	82	86	35.2	gy. m. and s.	do	
931	do	14 34 00	134 36 00	2,250	82	85	br. m.	Red clay	
932	do	14 17 00	134 46 00	2,330	91	83	35.2	br. m.	do	
933	Aug. 31	14 36 20	134 58 30	2,488	80	83	br. m.	do	
934	do	14 20 00	135 09 30	2,652	82	84	35.3	br. m.	do	
935	do	14 38 00	135 21 00	2,532	83	84	br. m.	do	
936	do	14 20 30	135 32 00	2,779	84	85	35.5	br. m.	Red clay	Do.
937	do	14 37 30	135 40 30	2,620	82	86	br. m.	do	
938	do	14 20 00	135 51 15	2,862	84	86	35.4	br. m.	do	
939	do	14 37 40	136 00 00	2,838	84	84	gy. m.	Diatom ooze	
940	Sept. 1	14 21 00	136 11 40	2,830	83	84	35.4	br. m.	do	Do.
941	do	14 39 15	136 20 30	2,748	82	83	br. m.	Red clay	
942	do	14 23 15	136 32 15	3,001	80	84	35.5	br. m.	do	Do.
943	do	14 40 30	136 41 00	2,838	89	85	yl. m.	Red clay	
944	do	14 24 00	136 53 00	2,559	87	85	35.2	br. m.	do	
945	do	14 41 00	137 01 40	2,877	83	85	br. m.	do	
946	do	14 23 00	137 13 00	2,751	83	84	35.2	br. m.	do	
947	Sept. 2	14 40 20	137 24 30	2,605	79	83	br. m.	do	
948	do	14 22 45	137 35 00	2,762	81	84	35.2	br. m.	do	Do.
949	do	14 38 00	137 49 00	2,725	83	84	br. c.	Red clay	
950	do	14 17 00	137 55 30	2,638	83	85	35.2	br. c.	Red clay	Do.

Abstract of the official record of soundings—Continued.

LUZON TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
951	Sept. 2	14 31 00	138 07 00	2,556	87	86		r	Red clay	Manganese concretions.
952	do	14 10 00	138 13 45	2,757	83	85	35.1	br. m.	do	
953	do	14 24 00	138 13 26	2,351	82	84		gvl. br. m. and s.	do	Manganese.
954	Sept. 3	14 04 15	138 31 15	2,646	82	83	35.2	br. m.	do	
955	do	14 18 20	138 46 30	2,793	82	85		br. m.	do	
956	do	13 57 30	138 52 00	2,763	84	86	35.2	br. m.	do	
957	do	14 12 00	139 08 30	2,473	84	86				No specimen.
958	do	13 54 20	139 11 15	2,298	86	86	35.1	br. m. and s.	Red clay	
959	do	14 13 00	139 26 45	3,180	83	85		wh. m.	Diatom ooze	
960	do	13 54 40	139 34 00	3,042	82	84	36	br. m. fn. bk. sp.	do	
961	Sept. 4	14 10 40	139 47 00	2,187	82	84				Do.
962	do	13 51 30	139 54 20	2,767	77	83	37			Do.
963	do	14 07 15	140 07 45	2,599	81	86		gvl.	Red clay	Pumice.
964	do	13 49 15	140 14 30	2,737	86	86		gvl.	do	
965	do	14 05 00	140 28 15	2,772	80	84		br. m.	do	
966	do	13 46 15	140 34 30	2,696	77	83		br. m.	do	
967	Sept. 5	14 01 30	140 49 00	2,706	82	84		br. m.	do	Radiolaria.
968	do	13 43 30	140 55 00	2,658	82	84		br. m.	do	
969	do	13 58 30	141 09 00	2,673	86	85		br. m.	do	
970	do	13 38 30	141 16 15	2,567	90	86		br. m.	do	
971	do	13 51 45	141 29 15	2,587	86	86		br. m.	do	
972	do	13 30 20	141 34 45	2,352	83	86	35			No specimen.
973	do	13 45 15	141 47 20	2,383	82	85		br. m.	Red clay	
974	Sept. 6	13 26 40	141 53 45	1,775	78	84	35.1	gy. m. and s.	Globigerina ooze.	
975	do	13 41 30	142 06 45	1,865	81	85		gy. fn. s.	do	
976	do	13 22 30	142 13 15	1,649	82	85		br. m. and s.	do	
977	do	13 38 30	142 24 30	1,755	87	86		gy. s. and m.	do	
978	do	13 21 00	142 32 30	1,380	88	86		wh. and bk. s.	do	
979	do	13 41 00	142 46 15	1,740	81	86				Do.
980	do	13 23 15	142 56 15	2,200	82	85		br. m.	Red clay	
981	Sept. 7	13 41 20	143 08 20	2,364	83	85	35.4	br. m.	do	
982	do	13 25 00	143 19 00	1,754	82	84		bk. s. and gvl.	Volcanic mud.	Many mang concretions.
983	do	13 43 45	143 31 45	1,882	80	84	35.1	bk gvl. bk. s.	do	Do.
984	do	13 26 45	143 42 30	1,751	85	85		brs. bk. and wh. s.	do	Do.
985	do	13 41 45	143 52 30	1,924	86	86	35.1	r	do	Do.
986	Sept. 9	13 34 30	144 31 30	1,411	80	86		bk. and gy. s. gy. m.	do	
987	do	13 37 00	144 14 30	1,889	87	86		fn. gy. m.		No specimen.
988	do	13 20 30	144 00 00	1,606	84	87		bk. s. gy. m.	Volcanic mud.	
989	do	13 17 00	144 19 45	1,927	83	87		gy. m.	Globigerina ooze.	Guam.

GUAM TO YOKOHAMA.

990	Sept. 9	13 28 30	144 36 15	859	83	87		fn. br. m.	Coralsand.	
991	do	13 38 30	144 35 00	1,143	81	86		fn. bk. and gy. s.	Globigerina ooze.	
992	do	13 40 30	144 34 45	1,013	83	86		r	do	Manganese.
993	do	13 45 45	144 34 00	1,970	83	86		r	do	
994	do	13 51 00	144 33 15	2,014	82	86		gy. m.	Volcanic mud.	Fine volcanic glass.
995	Sept. 10	14 00 45	144 31 45	2,091	83	85	35.2	vol. sand.	do	
996	do	14 02 45	144 31 30	2,005	83	85		br. m. and s.	do	
997	do	14 13 30	144 30 00	2,168	82	84		br. m. and s.	do	
998	do	14 15 40	144 29 45	2,158	82	84		fn. bk. s.	do	No specimen.
999	do	14 25 30	144 28 30	1,988	84	85		r	Volcanic mud.	Manganese.
1000	do	14 27 30	144 28 15	1,947	84	86	39	bk. and gy. s. and m.	do	
1001	do	14 37 30	144 27 00	2,005	86	86		bk. and gy. s. and m.	do	
1002	do	14 39 30	144 36 45	1,997	86	86		r		No specimen.
1003	do	14 49 30	144 25 20	2,233	81	86	36		Volcanic mud.	Do.
1004	do	14 51 40	144 25 00	2,214	83	86		r	do	Manganese.
1005	do	15 00 45	144 23 30	2,061	82	86		r	do	Do.
1006	do	15 02 45	144 23 15	1,847	83	86		bk. s. gy. m.	do	

Abstract of the official record of soundings—Continued.

GUAM TO YOKOHAMA—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur-face.	Bot-tom.			
1007	1899, Sept. 10	15 09 15	144 22 30	<i>Fath-oms.</i> 2,128	83	85	br. m. bk. s..	volcanic mud.	
1008do.....	15 20 15	144 20 45	1,985	81	85	br. m. bk. s..do.....	Pumice and manganese.
1009do.....	15 22 20	144 20 30	1,959	81	85	37	gvl.....do.....	
1010do.....	15 33 15	144 18 45	2,082	81	85	br. m. bk. s..do.....	
1011do.....	15 35 30	144 18 30	2,273	81	84	br. m.do.....	
1012do.....	15 46 30	144 16 45	1,932	81	84	35.1	br. m.do.....	
1013	Sept. 11	15 48 45	144 16 30	1,917	81	84	br. m.do.....	
1014do.....	15 59 45	144 14 45	1,987	81	84	br. m.do.....	
1015do.....	16 02 00	144 14 30	2,057	82	84	br. m.do.....	
1016do.....	16 13 00	144 12 40	2,375	83	85	br. m.do.....	
1017do.....	16 15 30	144 12 20	2,446	84	85	br. m. bk. s..do.....	
1018do.....	16 26 00	144 11 00	2,381	83	85	R. br. m.do.....	
1019do.....	16 28 00	144 10 45	2,211	87	86	35.5	br. m.do.....	
1020do.....	16 37 45	144 11 20	2,011	86	86	br. m. bk. s..do.....	
1021do.....	16 39 45	144 11 30	1,969	87	86	br. m. bk. s..do.....	
1022do.....	16 49 15	144 12 20	2,298	86	86	35.5	br. m.do.....	
1023do.....	16 51 00	144 12 15	2,392	83	86	br. m.do.....	
1024do.....	17 00 20	144 11 15	2,189	83	86	br. m.do.....	
1025do.....	17 02 15	144 11 00	2,271	84	85	br. m.do.....	
1026do.....	17 11 30	144 09 45	2,025	82	85	br. m.do.....	
1027do.....	17 13 30	144 09 30	2,314	83	85	br. m.do.....	
1028	Sept. 12	17 22 45	144 08 40	2,382	82	85	br. m. bk. s..do.....	
1029do.....	17 24 45	144 08 30	2,356	83	85	br. m.do.....	
1030do.....	17 34 15	144 07 45	2,112	83	85	br. m.do.....	
1031do.....	17 36 00	144 07 30	2,091	81	85	br. m. bk. s..do.....	
1032do.....	17 45 30	144 07 00	2,351	82	86	br. m. bk. s..do.....	
1033do.....	17 47 30	144 06 45	2,451	84	86	br. m. bk. s..do.....	
1034do.....	17 57 00	144 06 00	1,990	87	86	br. m. bk. s..do.....	
1035do.....	17 59 00	144 05 40	2,175	88	86	br. m.do.....	
1036do.....	18 08 30	144 04 45	2,155	86	86	br. m.do.....	
1037do.....	18 10 30	144 04 30	2,022	89	86	br. m. bk. s..do.....	
1038do.....	18 20 00	144 02 30	2,451	86	86	35.5	br. m. bk. s..do.....	
1039do.....	18 22 00	144 02 15	2,424	87	86	br. m. bk. s..do.....	
1040do.....	18 31 45	144 00 45	2,451	83	86	br. m.do.....	
1041do.....	18 33 45	144 00 30	2,433	84	85	R. br. m.do.....	Manganese.
1042do.....	18 43 30	143 59 15	2,225	83	84	br. m.do.....	
1043do.....	18 45 20	143 59 00	2,190	82	84do.....	No specimen.
1044	Sept. 13	18 55 00	143 57 30	2,303	83	84	35.7	br. m. bk. s..	Volcanic mud.	
1045do.....	18 57 00	143 57 15	2,330	83	83	br. m. bk. s..do.....	
1046do.....	19 06 00	143 56 30	2,220	82	83	br. m. bk. s..do.....	
1047do.....	19 08 00	143 56 15	2,133	82	84	br. m. bk. s..do.....	
1048do.....	19 17 30	143 55 00	1,967	85	85	36.8	br. m. bk. s..do.....	
1049do.....	19 19 30	143 54 45	1,964	84	86	br. m.do.....	Palagonite.
1050do.....	19 29 00	143 53 45	2,278	87	86do.....	
1051do.....	19 31 00	143 53 30	2,180	83	86	br. m.do.....	
1052do.....	19 40 15	143 52 45	2,146	81	85	br. m.do.....	No specimen.
1053do.....	19 42 15	143 52 30	2,151	82	85	br. m.	Volcanic mud.	
1054do.....	19 52 30	143 52 00	1,863	82	85	br. m.do.....	
1055do.....	19 54 45	143 52 00	2,028	82	85	35.7	br. m.do.....	
1056do.....	20 05 15	143 57 30	2,319	81	85	br. m. bk. s..do.....	
1057do.....	20 07 45	143 57 20	2,202	81	85	br. m.do.....	
1058do.....	20 18 20	143 51 00	1,930	82	85	35.6	br. m.do.....	
1059do.....	20 20 45	143 50 45	1,987	82	85	br. m.do.....	Do.
1060do.....	20 31 45	143 50 30	2,322	82	85	br. m.	Volcanic mud.	
1061	Sept. 14	20 34 30	143 50 30	2,181	82	85do.....	Do.
1062do.....	20 45 15	143 50 15	2,040	81	85	35.5	br. m.	Volcanic mud.	
1063do.....	20 47 45	143 50 15	1,884	82	84	br. m.do.....	
1064do.....	20 58 30	143 50 00	1,588	82	84	br. m.do.....	
1065do.....	21 01 00	143 50 00	1,321	82	84	br. m. bk. s..do.....	
1066do.....	21 06 30	143 50 00	1,815	84	85	br. m.do.....	
1067do.....	21 12 15	143 49 45	2,191	83	85	br. m. bk. s..do.....	
1068do.....	21 14 30	143 49 30	2,207	85	85	gy. m.do.....	
1069do.....	21 20 15	143 48 45	2,335	83	85	br. m.do.....	
1070do.....	21 30 00	143 47 30	1,714	83	85	35.1	bk. s. gy. m.do.....	
1071do.....	21 32 00	143 47 30	1,595	86	85	bk. s. gy. m.do.....	
1072do.....	21 37 15	143 46 30	1,470	85	86	R.do.....	Manganese.
1073do.....	21 42 45	143 45 30	1,208	85	86	S.do.....	
1074do.....	21 45 15	143 45 15	483	85	86	bk. s.	Volcanic sand.	

Abstract of the official record of soundings—Continued

GUAM TO YOKOHAMA—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
1075	Sept. 14	21 47 45	143 45 00	1,029	85	86		bk. s. m.	Volcanic mud.	
1076do....	21 53 00	143 44 00	1,530	84	86		bk. s. wh. sp.do....		
1077do....	21 58 15	143 43 00	1,547	84	86		gvl.do....		Manganese.
1078do....	22 00 45	143 42 45	1,465	84	86		gvl.do....		Do.
1079do....	22 06 20	143 41 45	1,547	83	85		fn. gvl.do....		Do.
1080do....	22 17 30	143 40 40	1,815	82	85		br. m. s.do....		Do.
1081do....	22 20 15	143 40 45	1,900	82	85		br. m. s.do....		Do.
1082	Sept. 15	22 31 30	143 40 45	2,083	82	85		bk. s. gy. m.do....		
1083do....	22 34 15	143 40 45	2,077	82	84		br. m.do....		
1084do....	22 45 30	143 40 45	2,313	83	84		br. m.do....		
1085do....	22 48 15	143 40 45	2,360	82	84		br. m.do....		
1086do....	22 59 30	143 41 15	2,677	83	84		br. m.do....		
1087do....	23 02 15	143 41 20	2,702	84	84		br. m. gvl.do....		
1088do....	23 13 30	143 41 45	2,952	83	85		br. m.do....		
1089do....	23 16 30	143 42 00	2,882	84	85		br. m.do....		
1090do....	23 27 30	143 41 20	2,725	89	86		gvl. s.do....		
1091do....	23 30 00	143 41 15	2,842	86	85		br. m. s.do....		
1092do....	23 40 20	143 40 30	3,189	85	85		br. m.do....		
1093do....	23 42 45	143 40 00	3,165	84	85		br. m.do....		
1094do....	23 52 30	143 37 45	3,595	83	84		br. m.do....	Red clay.	
1095do....	23 52 30	143 32 00	3,213	84	84		br. m.do....	Volcanic mud.	Volcanic glass.
1096	Sept. 16	23 52 30	143 26 00	2,998	83	84		br. m.do....	Red clay.	
1097do....	23 52 30	143 20 15	3,040	83	84		br. m.do....		
1098do....	23 57 40	143 19 45	3,259	82	84		br. m.do....	Volcanic mud.	Volcanic glass and radiolaria.
1099do....	23 59 40	143 19 40	3,251	84	84		gvl.do....		No specimen.
1100do....	23 59 40	143 14 15	2,483	84	84		br. m.do....	Volcanic mud.	
1101do....	24 04 40	143 13 45	2,855	85	85		br. m.do....		
1102do....	24 04 40	143 07 45	2,425	87	85	35	gvl.do....		Lumps of clay and manganese.
1103do....	24 09 40	143 07 30	2,294	87	85		br. m. and s.do....		
1104do....	24 14 30	143 06 45	1,904	87	85		R.do....		Volcanic glass.
1105do....	24 19 15	143 06 00	1,749	86	85	35.1	gy. s. m.do....		
1106do....	24 28 45	143 04 20	1,988	83	85		gvl.do....		Concretions of fine glass.
1107do....	24 30 45	143 04 15	2,190	84	85		gvl.do....		Do.
1108do....	24 39 30	143 04 15	2,645	83	85		br. m. bk. s.do....		
1109do....	24 41 30	143 04 15	2,662	82	85	35	br. m. and s.do....		
1110do....	24 50 30	143 04 20	2,870	83	85		br. m. bk. s.do....		
1111do....	24 52 30	143 04 20	2,788	83	84		gvl.do....		Concretions of glass.
1112	Sept. 17	25 01 30	143 04 30	2,564	81	83		br. m. bk. s.do....		
1113do....	25 04 30	143 04 30	2,555	83	84		br. m. s.do....		
1114do....	25 12 20	143 05 00	2,413	83	84	35.1	br. m. bk. s.do....		
1115do....	25 14 20	143 05 00	2,261	82	85		br. m. s.do....		
1116do....	25 22 45	143 05 30	2,186	84	85		br. m. s.do....		
1117do....	25 24 45	143 05 30	2,123	84	85		br. m. s.do....		
1118do....	25 33 30	143 06 00	1,805	82	85	35.5	br. m. bk. s.do....		
1119do....	25 35 30	143 06 00	1,654	82	85		br. m. s. gvl.do....		
1120do....	25 44 30	143 06 30	1,710	83	85		br. m. s.do....		
1121do....	25 46 30	143 06 45	1,887	82	86		br. m. s.do....		Brown glass and foraminifera.
1122do....	25 55 30	143 07 15	1,926	86	86		gy. m. s.do....		Do.
1123do....	25 57 30	143 07 15	1,877	86	86	35.1	gy. m. s.do....		
1124do....	26 06 45	143 07 45	1,229	82	85		R.do....		Manganese.
1125do....	26 08 45	143 08 00	1,251	83	85		gvl.do....		Foraminifera, manganese, and lumps of clay.
1126do....	26 12 45	143 08 00	972	82	85		gy. m. and s.do....		
1127do....	26 12 45	143 12 15	1,337	82	85	35.7	bk. s.do....		
1128do....	26 17 00	143 12 30	1,418	82	84		gy. m. and s.do....		Manganese and foraminifera.
....do....do....	26 21 30	143 12 45	1,505	82	84		gy. m. and s.do....		Do.
....do....	Sept. 18	26 30 30	143 13 15	2,304	83	84		gy. m. and s.do....		

Abstract of the official record of soundings—Continued.

GUAM TO YOKOHAMA—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1899.	° ' "	° ' "	Fathoms.	°	°	°			
1131	Sept. 18	26 32 30	143 13 15	2,351	82	84				No specimen.
1132	do	26 41 15	143 13 45	2,950	82	84	35.1	br. m. and s.	Volcanic mud.	
1133	do	26 43 45	143 13 45	2,800	82	84		br. m. and s.	do	
1134	do	26 47 45	143 14 00	2,879	83	85		br. m. and s.	do	
1135	do	26 52 20	143 13 00	3,421	86	85		br. m. and s.	do	
1136	do	26 52 20	143 07 00	3,132	86	85		bk. s. gy. m.	do	
1137	do	26 52 15	143 00 20	2,250	89	86		gy. m. and s.	do	
1138	do	26 48 00	143 02 15	1,837	87	86	35	gy. s. and m.	do	Manganese.
1139	do	26 43 30	143 04 15	2,101	85	86		gy. s. and m.	do	No specimen.
1140	do	26 43 30	142 57 45	1,835	88	86		gy. s. and m.	do	
1141	do	26 55 30	142 58 30	2,278	82	86		gy. s. and m.	Volcanic mud.	
1142	do	27 04 40	142 57 00	2,682	83	86		br. s. and m.	do	Radiolaria and diatoms.
1143	do	27 07 40	142 56 30	2,591	82	86	35.1	br. s. and m.	do	
1144	Sept. 19	27 17 30	142 54 45	2,543	82	84		br. m. and s.	do	
1145	do	27 19 30	142 54 20	2,119	81	84		bk. and gy. s. and m.	do	
1146	do	27 24 30	142 53 20	2,251	80	84		bk. and gy. s. and m.	do	
1147	do	27 29 30	142 52 20	1,856	83	84		gy. s. and m.	Globigerina ooze.	
1148	do	27 34 30	142 51 30	2,181	81	84		bk. and gy. s. and m.	Volcanic mud.	
1149	do	27 39 15	142 50 45	2,106	84	84		gy. C. bk. s.	do	
1150	do	27 48 45	142 49 45	1,746	82	84		gy. m. and s.	Globigerina ooze.	
1151	do	27 57 00	142 49 30	1,686	83	84		gy. m. and s.	Volcanic mud.	Many foraminifera. Do. Do.
1152	do	28 01 00	142 48 45	2,041	83	84	35.2	gy. C.	do	
1153	do	28 03 00	142 48 15	1,932	82	84		gy. m. and s.	do	
1154	do	28 12 20	142 43 30	1,602	82	85		gy. s.	do	
1155	do	28 14 30	142 42 30	1,632	82	85		gy. m. and s.	do	
1156	do	28 23 45	142 37 45	1,660	83	85		gy. m. and s.	do	
1157	do	28 25 15	142 36 45	1,617	83	85		gy. m. and s.	do	
1158	do	28 35 20	142 33 45	1,584	82	85		gy. m. and s.	do	
1159	do	28 37 20	142 32 15	1,515	82	85		gy. m. and s.	Volcanic mud.	No specimen. Foraminifera. Do.
1160	do	28 46 45	142 28 15	1,907	82	85		gy. m. and s.	do	
1161	do	28 49 00	142 27 20	1,994	82	85	37	gy. m. and s.	do	
1162	Sept. 20	28 58 40	142 23 30	2,095	82	85		gy. m. and s.	Globigerina ooze.	
1163	do	29 00 45	142 22 30	2,049	82	83		gy. m. and s.	do	
1164	do	29 10 30	142 18 30	2,384	83	83		gy. m. and s.	Volcanic mud.	
1165	do	29 12 30	142 17 40	2,387	82	84		gy. m. and s.	do	
1166	do	29 22 20	142 13 30	2,552	86	85	35	br. m. and s.	do	
1167	do	29 24 20	142 12 30	2,596	86	85		br. m. and s.	do	
1168	do	29 34 00	142 08 30	2,933	85	85		br. m. bk. s.	do	
1169	do	29 36 00	142 07 30	2,927	90	86		C.	do	No specimen.
1170	do	29 45 00	142 02 15	2,912	86	86	35	gy. m. and c.	Volcanic mud.	
1171	do	29 46 45	142 01 15	2,826	88	86		br. m. and s.	do	
1172	do	29 55 45	141 55 30	2,621	84	86		br. m. and s.	do	
1173	do	29 57 40	141 54 30	2,655	82	85		bk. s.	do	
1174	do	30 06 00	141 48 30	2,490	82	85		br. m. bk. s.	do	
1175	do	30 07 45	141 47 30	2,384	84	85		br. m. and s.	do	
1176	Sept. 21	30 16 30	141 41 15	2,089	82	84		br. m. and s.	do	
1177	do	30 18 15	141 40 00	1,987	82	84	35.1	gy. m. and s.	do	
1178	do	30 26 45	141 33 10	1,685	82	84		gy. m. and s.	do	
1179	do	30 28 30	141 32 15	1,652	82	84		bk. s.	do	
1180	do	30 36 45	141 25 30	1,617	81	85	35.2	gy. m. bk. s.	do	
1181	do	30 38 30	141 24 15	1,590	82	85		gy. m. bk. s.	do	
1182	do	30 46 45	141 17 00	1,548	83	85		gvl.	do	Concretions of volcanic glass. Manganese.
1183	do	30 48 30	141 16 00	1,454	82	85		gy. m. bk. s.	do	
1184	do	30 57 15	141 10 45	1,542	82	85		br. m. and s.	do	
1185	do	30 59 00	141 09 30	1,491	82	85	35.2	gy. m. and s.	do	
1186	do	31 08 15	141 06 45	1,842	82	85		gy. m. and s.	do	
1187	do	31 10 00	141 06 30	1,815	86	85		gy. m. and s.	do	

Abstract of the official record of soundings—Continued.

GUAM TO YOKOHAMA—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Fathoms.	Air.	Surface.	Bottom.		
1188	1899. Sept. 21	° ' "	° ' "			°	°	°		
		31 18 15	141 03 30	1,595	80	85	35.2	gy. m. gy. and bk. s.	Volcanic mud.	
1189do	31 20 00	141 02 45	1,562	80	84	gy. m. gy. and bk. s.do	
1190do	31 27 40	141 00 30	1,543	79	84	gy. m. and s.do	
1191do	31 29 15	141 00 00	1,551	79	84	gy. m. and s.do	
1192do	31 37 45	140 57 15	1,545	78	83	gy. m. and s.do	
1193do	31 39 20	140 56 30	1,512	77	82	gy. m. and s.do	
1194	Sept. 22	31 48 00	141 03 45	1,547	76	80	gy. m. and s.do	
1195do	31 57 40	141 00 45	1,431	77	82	gy. m. and s.do	
1196do	31 59 15	141 00 00	1,431	77	82	gy. m. and s.do	
1197do	32 10 30	140 56 30	1,698	76	81	gy. m. and s.do	
1198do	32 13 15	140 55 45	1,612	76	81	gy. m. and s.do	
1199do	32 24 15	140 52 00	1,497	76	80	gy. and br. m. and s.do	
1200do	32 27 15	140 51 15	1,460	76	80	gy. m. gy. and bk. s.do	
1201do	32 39 30	140 47 30	1,268	76	81	gy. m. gy. and bk. s.do	
1202do	32 42 00	140 46 45	1,199	75	81	gy. m. gy. and bk. s.do	
1203	Sept. 23	32 53 30	140 43 30	908	72	79	gy. m. gy. and bk. s.do	
1204do	32 56 15	140 43 00	846	72	79	bk. s.do	Concretions of sand and manganese.
1205do	33 07 30	140 39 40	737	74	79	bk. s.do	
1206do	33 10 30	140 39 00	737	74	78	bk. s. m.do	
1207do	33 22 00	140 35 45	665	75	79	bk. s. m.	Blue mud	
1208do	33 25 00	140 35 00	665	74	79	bk. s. m.do	
1209do	33 36 15	140 32 15	660	74	80	bk. s. gy. m.do	
1210do	33 38 45	140 31 15	688	74	80	bk. gy. s.do	
1211do	33 50 00	140 28 30	777	74	80	bk. gy. s.do	
1212do	33 52 00	140 27 40	815	74	80	gvl.do	
1213do	34 01 00	140 24 15	808	73	80	bk. gy. s.do	Do.
1214do	34 03 15	140 24 00	857	73	80	bk. s. gvl.do	
1215do	34 12 00	140 16 15	901	73	80	bk. s. gn. m.do	
1216do	34 14 00	140 14 00	920	73	80	36.6	bk. s. gn. m.do	
1217do	34 21 45	140 04 00	934	72	80	36.2	gn. m.	Green mud	
1218do	34 23 00	140 01 45	932	73	77	gn. m.do	
1219do	34 25 00	139 57 45	879	73	77	gn. m.do	
1220do	34 26 45	139 53 30	786	73	77	36	gn. m. br. s.do	
1221do	34 28 30	139 49 30	781	71	78	gn. m. bk. s.do	
1222do	34 30 00	139 45 15	726	71	78	gn. m. bk. s.do	
1223do	34 31 45	139 41 00	676	71	78	gn. m. bk. br. s.do	
1224	Sept. 24	34 33 00	139 36 30	660	70	78	gn. m. bk. br. s.do	
1225do	34 41 00	139 33 15	805	69	78	41.1	bk. s. gn. m.do	
1226do	34 44 30	139 32 00	977	69	77	bk. s. gn. m.do	
1227do	34 47 20	139 31 00	812	69	77	bk. s. gn. m.do	
1228do	34 50 00	139 26 00	792	69	77	bk. s. gn. m.do	
1229do	34 50 45	139 24 45	650	69	77	gvl.	Manganese nodules.	Fine sediment, washed out.
1230do	34 51 00	139 24 00	689	69	77	gn. m. bk. s.do	
1231do	34 51 30	139 23 00	807	69	77	gn. m. bk. s.do	
1232do	34 54 00	139 20 20	905	69	77	gn. m. bk. s.do	
1233do	34 57 45	139 19 40	807	68	75	gn. m. bk. and br. s.do	
1234do	35 00 30	139 21 45	805	68	75	gn. m. bk. s.do	
1235do	35 03 40	139 24 00	720	69	75	gn. m. bk. s.do	
1236do	35 06 30	139 25 15	652	69	75	gn. m. bk. s.do	
1237do	35 07 30	139 26 00	613	69	75	gn. m. bk. s.do	Yokohama.

YOKOHAMA TO GUAM.

1238	Oct. 10	34 51 30	139 37 30	272	68	73	bk. s. and gvl.	Green mud	
1239do	34 48 45	139 35 30	1,005	69	74	bk. m. and s.do	
1240do	34 47 15	139 40 30	612	69	74	bk. and gy. m. and s.do	
1241do	34 43 15	139 38 00	1,115	69	74	bk. and gy. m. and s.do	

OCEANOGRAPHY OF THE PACIFIC.

47

Abstract of the official record of soundings—Continued.

YOKOHAMA TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur-face.	Bot-tom.			
	1899.	° ' "	° ' "	Fath-oms.	°	°	°			
1242	Oct. 10	34 44 30	139 49 30	797	72	74	bk. and gy. m. and s.	Green mud.	
1243do	34 37 15	139 50 00	1,277	69	74	bk. s.do	
1244do	34 39 15	140 00 00	1,363	69	74	bk. and gy. m. and s.do	
1245do	34 29 45	139 58 45	1,098	69	74	gvl.do	
1246	Oct. 11	34 20 20	139 57 30	1,299	70	70	bk. and gy. m. and s.do	
1247do	34 29 45	140 23 30	1,814	70	75	No specimen.
1248do	34 21 15	140 21 30	1,742	70	75	bk. s.do	Do.
1249do	34 13 00	140 19 30	1,323	69	75	bk. and gy. m. and s.	Blue mud	
1250do	34 06 00	140 10 00	733	70	75	bk. s.do	
1251do	34 08 15	140 33 00	1,270	72	78	gy. m.do	
1252do	33 50 30	140 20 00	745	70	77	37	gy. and bk. m. and s.do	
1253do	33 47 45	140 38 15	1,194	70	77	gy. m.do	
1254do	33 32 40	140 20 00	264	69	76	fn. gvl.	Coral sand.	Pteropods and foraminifera.
1255do	33 30 00	140 25 45	439	68	76	fn. gvl.	Blue mud	
1256do	33 27 45	140 30 45	600	70	76	fn. bk. s.do	
1257	Oct. 12	33 23 00	140 41 30	812	69	76	gy. m.do	
1258do	33 05 45	140 24 00	454	67	76	43.7	gvl.do	Manganese.
1259do	32 57 30	140 45 00	964	70	79	gy. m. bk. s.do	
1260do	32 56 15	140 50 00	1,094	74	80	gy. m.	Volcanic mud.	
1261do	32 47 45	140 35 45	920	70	80	gy. m. bk. s.do	
1262do	32 39 30	140 57 30	1,428	72	78	br. m. bk. s.do	
1263do	32 28 45	140 37 45	1,246	72	78	35.4	br. m. bk. s.do	Manganese and volcanic glass
1264do	32 22 00	141 02 30	2,080	73	79	bk. s.do	
1265do	32 10 30	140 44 00	1,444	71	78	35.2	gy. m.do	
1266	Oct. 13	32 05 00	141 07 30	1,730	73	77	gy. m.do	
1267do	31 54 00	140 49 00	1,461	71	77	35.3	gy. m.do	
1268do	31 48 00	141 11 15	1,651	74	77	gy. m.do	
1269do	31 36 20	140 53 30	1,622	75	80	35	gy. m. bk. s.do	
1270do	31 29 45	141 16 00	1,915	75	80	gy. m.do	
1271do	31 17 30	140 58 00	1,557	77	80	gy. m.do	
1272do	31 12 30	141 18 30	2,165	71	79	gy. m. bk. s.do	
1273do	31 00 30	140 58 30	1,463	75	78	35.1	gy. m. bk. s.do	
1274	Oct. 14	30 56 30	141 18 40	1,600	73	78	gy. m. bk. s.do	
1275do	30 41 45	141 01 45	1,620	73	78	gvl.do	
1276do	30 42 00	141 23 00	1,807	74	79	35	gvl. gy. m.do	
1277do	30 29 30	141 12 30	1,857	77	80	gvl. crs. bk. s.do	
1278do	30 28 30	141 41 30	2,266	75	80	35	gy. m. bk. s.do	
1279do	30 51 00	141 29 00	2,175	75	80	gvl. bk. s.do	
1280do	30 19 00	141 24 30	2,215	79	81	35	bk. s.do	
1281	Oct. 15	30 19 30	141 48 00	2,558	78	80	gy. m. bk. s.do	
1282do	30 07 00	141 37 00	2,299	80	81	35	gy. m. bk. s.do	
1283do	30 05 20	142 02 30	2,767	84	81	gy. m. bk. s.do	
1284do	29 50 45	141 29 00	2,588	79	81	gy. m. bk. s.do	
1285do	29 46 20	142 13 00	3,576	80	81	35.2	gy. m. bk. s.do	
1286	Oct. 17	29 23 30	141 45 30	1,606	77	80do	Typhoon; no specimen.
1287do	29 24 00	141 54 45	1,606	78	80	35	gy. m. bk. s.	Volcanic mud.	
1288do	29 25 00	142 03 15	1,755	78	80	wh. and bk. s.	Globigerina ooze.	
1289do	29 25 45	142 13 00	2,651	74	80	br. m.	Volcanic mud.	
1290do	29 21 15	142 34 00	4,350	76	81	br. m.do	
1291do	29 32 20	142 21 30	4,212	79	80	br. m.do	
1292do	29 36 00	142 00 00	2,371	74	80	35	br. m. bk. and wh. s.do	
1293do	29 37 15	141 50 45	2,141	73	80	br. m.do	
1294	Oct. 18	29 17 30	142 04 30	1,417	74	79	wh. gy. bk. s.	Globigerina ooze.	
1295do	29 10 30	141 57 30	1,415	74	79	gy. bk. s. and gy. m.do	
1296do	29 09 00	142 08 40	1,758	73	79	gy. m.do	
1297do	29 00 45	142 12 00	1,954	74	81	gy. m.	Volcanic mud.	

Abstract of the official record of soundings—Continued.

YOKOHAMA TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1899.	° ' "	° ' "	Fathoms.	°	°	°			
1298	Oct. 18	28 53 30	142 05 40	1,711	76	81	gy. m. and gy. wh. s.	Globigerina ooze.	
1299	do	28 50 00	142 17 00	1,817	76	82	gy. m. and gy. wh. s.	do	
1300	do	28 40 40	142 21 00	1,529	80	81	gy. m.	do	
1301	do	28 33 00	142 14 00	1,088	77	81	35.9	gy. m. s.	do	
1302	do	28 31 00	142 25 00	1,331	75	81	gy. m. s.	do	
1303	do	28 21 15	142 28 15	1,410	75	81	35.1	gy. m. s.	do	
1304	do	28 13 00	142 20 30	847	76	81	gy. m. bk. s.	do	Much manganese and volcanic sand.
1305	do	28 10 30	142 31 00	1,289	75	80	gy. m.	do	
1306	do	28 00 00	142 34 00	1,208	75	81	gy. m. bk. s.	do	
1307	Oct. 19	27 52 40	142 25 00	616	76	81	38.3	gy. m. bk. s.	do	
1308	do	27 49 00	142 34 30	1,040	75	80	bk. and gy. s. and m.	do	
1309	do	27 38 30	142 34 30	891	77	80	bk. and gy. s.	do	Do.
1310	do	27 33 30	142 23 30	518	78	80	bk. and gy. s.	do	Do.
1311	do	27 54 40	142 42 15	1,503	80	82	gy. m.	do	
1312	do	27 40 30	142 43 15	1,552	76	82	gy. m.	do	
1313	do	27 30 30	142 44 15	1,716	79	82	gy. m. bk. s.	Volcanic mud.	
1314	do	27 22 45	142 45 30	1,660	81	82	gy. m. bk. s.	do	
1315	do	27 20 45	142 45 40	1,494	81	82	gy. m. bk. s.	do	Manganese and foraminifera.
1316	do	27 19 00	142 45 45	1,649	80	82	35	gy. m. bk. s.	do	Do.
1317	do	27 18 30	142 36 30	1,453	77	82	gy. m. bk. s.	do	Do.
1318	do	27 13 30	142 46 30	1,210	76	81	gy. m. bk. s.	do	Do.
1319	do	27 08 15	142 47 30	2,167	78	81	35	br. m. bk. s.	do	
1320	do	27 02 45	142 48 30	2,048	78	81	gy. m. bk. s.	do	Foraminifera.
1321	Oct. 20	26 56 00	142 41 00	1,618	78	81	35	gy. m. bk. s.	do	Do
1322	do	26 49 45	142 51 40	2,142	78	81	gy. m. bk. s.	do	
1323	do	26 38 00	142 53 00	1,583	78	81	bk. and gy. s.	do	
1324	do	26 31 00	142 44 15	1,915	79	81	35	gy. m.	do	
1325	do	26 25 00	142 55 00	847	80	82	br. m. bk. s.	do	
1326	do	26 30 30	143 02 15	865	79	82	bk. and wh. s.	do	
1327	do	26 15 30	142 56 00	1,591	80	83	35	s. gy. m.	do	
1328	do	26 20 30	142 56 00	871	79	83	bk. s. gvl.	do	Foraminifera and volcanic glass.
1329	do	26 22 00	142 50 40	1,709	82	83	gy. m. bk. s.	do	Manganese iron.
1330	do	26 23 30	142 45 00	1,543	81	83	gvl.	do	
1331	do	26 25 45	142 39 45	1,257	81	82	35.6	gy. m.	do	
1332	do	26 38 00	142 50 00	1,807	80	82	gy. bk. s.	do	
1333	do	26 36 00	142 37 00	1,186	79	82	35.1	gy. m.	do	
1334	do	26 16 45	142 41 00	1,334	78	82	gy. m. bk. s.	do	
1335	do	26 14 00	142 51 15	1,525	79	82	35	s. gvl.	do	
1336	Oct. 21	26 08 20	142 59 00	1,521	79	81	gy. bk. s.	do	
1337	do	26 02 45	143 06 40	1,485	80	81	35.6	br. wh. s.	Volcanic mud.	No specimen.
1338	do	25 57 30	142 59 00	1,424	79	82	bl. br. s. gy. m.	do	
1339	do	25 52 30	142 51 20	1,801	78	82	gy. m. bk. s.	do	
1340	do	25 46 15	143 01 15	2,000	81	81	35	gvl. bk. s.	do	Manganese nodules.
1341	do	25 37 15	143 57 00	1,602	80	82	br. bk. s.	do	Brown glass.
1342	do	25 32 20	142 47 45	1,710	81	83	35	gy. m. bk. s.	do	
1343	do	25 28 00	142 57 40	1,748	84	83	bk. s. gvl.	do	Manganese glass, and foraminifera.
1344	do	25 18 00	142 57 30	1,995	87	83	gy. m. bk. s.	do	
1345	do	25 13 20	142 47 30	1,449	81	83	35.5	bk. s. gvl.	do	
1346	do	25 07 30	142 57 00	1,755	79	83	35	gy. bk. s. gy. m.	do	
1347	do	24 57 00	142 56 30	2,482	80	83	35.2	gy. m. bk. s.	do	
1348	do	24 59 45	142 52 00	2,207	80	83	35	br. m. bk. s.	do	
1349	do	25 02 30	142 47 30	1,624	80	83	35	gvl.	do	Manganese.
1350	Oct. 22	24 52 00	142 47 00	2,249	79	81	35	gy. m. bk. s.	do	
1351	do	24 49 00	142 52 00	2,427	79	81	35	gy. m. bk. s.	do	

Abstract of the official record of soundings—Continued.

YOKOHAMA TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
1352	1899. Oct. 22	24 46 00	142 56 30	Fath- oms. 2,355	80	81	35	br. m. bk. s.	Volcanic mud.	
1353	...do...	24 40 40	142 56 15	2,065	80	82	...	gy. m. bk. s.	...do...	
1354	...do...	24 35 30	142 56 00	2,095	81	83	35	gy. m. bk. s.	...do...	
1355	...do...	24 32 40	142 51 30	1,843	81	83	35	gy. m. bk. s.	...do...	
1356	...do...	24 30 00	142 47 00	1,749	83	80	35	gy. m. bk. s.	...do...	
1357	...do...	24 24 30	142 56 00	1,735	82	83	35	gy. m. bk. s.	...do...	
1358	...do...	24 17 00	142 58 30	1,774	85	84	35.1	gy. m. bk. s.	...do...	
1359	...do...	24 10 30	142 50 40	1,737	84	85	...	gvl	...do...	Fibrous glass.
1360	...do...	24 07 20	143 00 30	2,368	88	85	35	gy. m. bk. s.	...do...	
1361	...do...	24 01 15	142 52 30	2,373	79	84	35	br. m. bk. s.	...do...	
1362	...do...	23 57 45	143 03 30	2,673	81	84	35	br. m. bk. s.	...do...	
1363	...do...	23 54 20	143 14 15	2,599	80	83	...	br. m. bk. s.	...do...	
1364	Oct. 23	23 27 45	143 06 4	2,744	80	83	35	br. m. bk. s.	...do...	
1365	...do...	23 41 45	142 59 15	2,272	81	83	35	br. m. bk. s.	...do...	
1366	...do...	23 38 00	143 10 00	2,368	81	84	36	br. m. bk. s.	...do...	
1367	...do...	23 34 30	143 21 30	2,609	84	84	35.1	br. m. bk. s.	...do...	
1368	...do...	23 28 00	143 13 30	2,323	84	84	...	bk. s. gy. m.	...do...	
1369	...do...	23 21 45	143 06 00	1,952	81	84	35	bk. s. gy. m.	...do...	
1370	...do...	23 18 15	143 16 40	2,588	85	84	35.1	br. m.	...do...	
1371	...do...	23 14 20	143 27 30	2,961	79	84	35	br. m.	...do...	
1372	...do...	23 08 15	143 20 00	2,582	81	84	35	br. m.	...do...	
1373	...do...	23 02 00	143 12 45	2,478	80	84	35	br. m. bk. s.	...do...	
1374	Oct. 24	22 57 30	143 22 45	2,548	81	84	35	br. m. bk. s.	...do...	
1375	...do...	22 52 45	143 33 00	2,483	81	84	35	br. m. bk. s.	...do...	
1376	...do...	22 48 15	143 24 30	2,415	81	84	35	br. m.	...do...	
1377	...do...	22 43 30	143 16 30	2,127	81	84	35	gy. m.	...do...	
1378	...do...	22 38 45	143 26 30	2,122	81	85	...	gy. m. bk. s.	...do...	
1379	...do...	22 34 15	143 36 30	2,131	86	85	35	gy. m. bk. s.	...do...	
1380	...do...	22 29 45	143 28 00	2,023	85	85	35	gy. m. bk. s.	...do...	
1381	...do...	22 25 00	143 18 40	1,099	86	85	35	bk. gy. s. gvl.	...do...	
1382	...do...	22 15 00	143 22 20	1,297	84	85	35.3	bk. gy. s.	...do...	
1383	...do...	22 19 00	143 26 00	1,388	83	85	35.6	bk. gy. s. gy.	...do...	
1384	...do...	22 14 45	143 31 45	1,532	82	85	35.3	br. m. bk. s.	...do...	
1385	...do...	22 11 30	143 26 00	1,236	82	85	36	br. m. bk. s.	...do...	
1386	...do...	22 08 15	143 20 15	802	82	84	37	bk. gy. s.	...do...	
1387	...do...	22 02 30	143 26 30	1,197	82	84	...	bk. s.	...do...	
1388	Oct. 25	21 48 36	143 40 45	1,668	81	83	35.5do...	No specimen.
1389	...do...	21 45 45	143 41 15	1,653	82	88	...	br. m. bk. s.	Volcanic mud.	
1390	...do...	21 42 30	143 41 30	1,801	81	83	35.2	gy. m. bk. s.	...do...	
1391	...do...	21 39 15	143 42 00	1,849	81	80	35.3	gy. m. bk. s.	...do...	
1392	...do...	21 36 00	143 42 30	1,615	82	84	35.2	gy. m. bk. s.	...do...	
1393	...do...	21 42 00	143 49 00	1,971	82	84	...	bk. s.	...do...	
1394	...do...	21 48 00	143 55 00	1,460	82	84	...	br. s. and m.	...do...	
1395	...do...	21 50 00	143 43 00	1,248	88	85	36.5	bk. s.	...do...	
1396	...do...	21 45 15	143 42 45	1,046	85	85	...	r	...do...	Manganese.
1397	...do...	21 58 15	143 33 00	1,053	82	85	36	bk. s. gvl.	...do...	Brown glass.
1398	...do...	21 57 30	143 38 00	1,392	87	85	35.5	bk. s.	...do...	
1399	...do...	21 54 00	143 34 00	1,215	84	85	36	bk. s.	...do...	
1400	...do...	21 49 40	143 29 15	1,374	83	85	35.5	bk. s.	...do...	
1401	...do...	21 46 15	143 34 00	1,594	81	85	35.6	bk. s.	...do...	
1402	...do...	21 43 30	143 29 00	1,715	83	84	35.2	bk. s.	...do...	
1403	...do...	21 40 00	143 24 00	1,820	83	84	...	bk. s.	...do...	
1404	Oct. 26	21 37 00	143 33 30	1,489	81	84	87	bk. s. gvl.	...do...	
1405	...do...	21 29 30	143 26 00	1,962	81	84	...	br. m. bk. s.	...do...	
1406	...do...	21 26 45	143 35 00	1,692	81	85	...	r	...do...	Manganese concretions.
1407	...do...	21 19 30	143 27 00	1,865	81	85	35.3	br. m. bk. s.	...do...	
1408	...do...	21 17 00	143 36 30	1,620	81	85	35.5	br. m. bk. s.	...do...	
1409	...do...	21 14 30	143 46 15	2,209	86	85	35.4	br. m. bk. s.	...do...	
1410	...do...	21 28 15	143 56 30	1,898	83	85	35.5	br. m. bk. s.	...do...	
1411	...do...	21 38 00	143 57 15	1,956	82	85	...	br. m. bk. s.	...do...	
1412	...do...	21 57 30	143 52 00	1,144	81	85	36	br. m. and s.	...do...	
1413	...do...	21 52 15	143 30 30	838	82	85	36.7	bk. s.	...do...	
1414	...do...	21 47 00	143 49 00	1,714	82	84	...	bk. s. br. m.	...do...	
1415	Oct. 27	21 46 30	144 02 00	1,300	81	84	35.5	br. m. and s.	...do...	
1416	...do...	21 34 45	144 03 00	1,912	80	84	...	bk. s.	...do...	
1417	...do...	21 15 15	143 56 30	1,691	81	84	35.2	br. m. bk. s.	...do...	

Abstract of the official record of soundings—Continued.

YOKOHAMA TO GUAM—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
1418	1899. Oct. 27	21 05 30	143 36 45	Fath- oms. 2,143	81	84		br. m. bk. s.	Volcanic mud.	
1419	...do...	20 53 00	143 55 15	1,874	84	85	35.3	br. m. and s.	...do...	
1420	...do...	20 46 30	143 39 40	2,095	85	85		br. m. bk. s.	...do...	
1421	...do...	20 39 45	143 59 30	1,889	82	84		br. m. bk. s.	...do...	
1422	...do...	20 31 00	143 43 00	2,250	83	84	35.5	br. m. bk. s.	...do...	
1423	Oct. 28	20 24 15	144 03 45	2,139	82	84		br. m. bk. s.	...do...	
1424	...do...	20 16 00	143 47 15	1,831	82	84		br. m. bk. s.	...do...	
1425	...do...	20 09 30	144 08 00	1,833	82	84	35.5	br. m. bk. s.	...do...	
1426	...do...	19 59 45	143 48 00	2,151	84	84		br. m. bk. s.	...do...	
1427	...do...	19 51 45	144 03 15	2,472	85	85	35.6	br. m. bk. s.	...do...	
1428	...do...	19 39 00	143 44 00	1,981	88	85	35.3	br. m. bk. s.	...do...	
1429	...do...	19 28 40	144 03 40	1,972	82	85	35.3	br. m. bk. s.	...do...	
1430	...do...	19 15 30	143 46 30	2,433	82	85	35.3	br. m. bk. s.	...do...	
1431	Oct. 29	19 06 15	144 06 00	2,308	82	84	35.4	br. m. bk. s.	...do...	
1432	...do...	18 53 30	143 48 15	1,906	82	84	35.1	bk. s.	...do...	
1433	...do...	18 44 15	144 05 15	2,169	83	84		br. m. bk. s.	...do...	
1434	...do...	18 33 00	143 49 45	2,202	86	85	35.2	br. m. bk. s.	...do...	
1435	...do...	18 25 45	144 12 00	2,349	81	85	35.6	br. m. bk. s.	...do...	
1436	...do...	18 13 20	143 54 00	2,265	84	84	35.2	br. m. bk. s.	...do...	
1437	...do...	18 05 00	144 13 30	2,127	80	84	35.2	br. m. bk. s.	...do...	
1438	Oct. 30	17 53 00	143 56 00	2,114	82	84		br. m. bk. s.	...do...	
1439	...do...	17 44 30	144 15 30	1,901	82	84	35.3	br. m. bk. s.	...do...	
1440	...do...	17 32 45	143 58 00	1,737	83	84	35.1	br. m. bk. s.	...do...	
1441	...do...	17 24 00	144 17 20	2,036	81	85		br. m. bk. s.	...do...	
1442	...do...	17 11 30	144 00 00	2,002	91	85	35	bk. br. s. br. m.	...do...	
1443	...do...	17 03 00	144 18 00	2,329	82	84	35.4	bk. br. s. br. m.	...do...	
1444	...do...	16 51 30	143 59 45	2,175	83	84	35.1	bk. br. s. br. m.	...do...	
1445	...do...	16 42 45	144 18 00	2,043	82	84		bk. br. s. br. m.	...do...	
1446	Oct. 31	16 31 00	143 59 30	2,113	82	84	35	bk. br. s. br. m.	...do...	
1447	...do...	16 23 00	144 17 30	2,447	83	84	35.3	bk. br. s. br. m.	...do...	
1448	...do...	16 11 00	143 58 30	2,084	81	85		bk. br. s. br. m.	...do...	
1449	...do...	16 01 20	144 20 20	2,365	82	85	35.1	bk. br. s. br. m.	...do...	
1450	...do...	15 51 30	144 04 15	2,401	85	85	35.1	br. m. bk. s.	...do...	
1451	...do...	15 45 00	144 26 20	1,801	80	84	35.2	br. m. bk. s.	...do...	
1452	...do...	15 34 45	144 12 00	2,116	79	84	35	br. m. bk. s.	...do...	
1453	Nov. 1	15 20 15	144 32 45	2,164	79	84	35	br. m. bk. s.	...do...	
1454	...do...	15 16 30	144 16 45	2,221	81	84		br. m. bk. s.	...do...	
1455	...do...	15 10 30	144 36 15	2,000	84	85		br. m. bk. s.	...do...	
1456	...do...	14 57 30	144 18 30	2,245	86	86		br. m. bk. s.	...do...	
1457	...do...	14 50 45	144 37 00	1,970	85	85	35.2	br. m. bk. s.	...do...	
1458	...do...	14 37 45	144 17 15	2,339	86	85		gy. m. and s.	...do...	
1459	...do...	14 30 40	144 36 20	1,981	82	84	35	gy. m. bk. s.	...do...	
1460	Nov. 2	14 17 30	144 19 30	2,053	83	84	35	c	...do...	Fragments of volcanic glass.
1461	...do...	14 10 00	144 39 00	1,946	82	84	35	gvl. br. m.	...do...	
1462	...do...	13 57 20	144 23 00	2,111	81	84	35	br. m. br. bk. s.	...do...	Volcanic glass.
1463	...do...	13 49 00	144 43 15	951	76	84		br. m. br. bk. s.	Globigerina ooze.	
1464	...do...	13 47 40	144 38 00	891	80	86		gy. m. bk. s.	...do...	
1465	...do...	13 47 00	144 35 30	993	83	85		r	...do...	San Luis d' Apra, Guam.

GUAM TO MIDWAY ISLANDS.

1466	Nov. 12	13 26 30	144 36 37	234	78	co. s. and m.	Coral mud.	No specimen.
1467	...do...	13 39 20	144 41 20	518	79	84	...	co. s. and m.	Globigerina ooze.	
1468	...do...	13 46 15	144 25 45	1,731	87	84	...	co. s. and m.	...do...	
1469	...do...	13 20 20	144 28 15	1,017	82	84	...	co. s. and m.	...do...	
1470	...do...	13 15 15	144 26 45	679	81	84	40.5	co. s. and m.	Globigerina ooze.	
1471	...do...	13 12 15	144 22 30	503	80	84	39.8	co. s. and m.	...do...	

Abstract of the official record of soundings—Continued.

GUAM TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	<i>Fath- oms.</i>	°	°	°			
1472	Nov. 12	13 17 00	144 32 00	1,000	81	84	co. s. gy. s. . .	Globigerina ooze.	Gren sand Mollusks.
1473	do	13 05 15	144 34 30	716	81	84	co. s. gy. s.	
1474	do	12 58 30	144 42 45	1,074	82	84	37.4	bk. s. g.	No specimen.
1475	Nov. 13	12 58 30	144 53 30	1,817	81	84	co. s. and m. . .	Globigerina ooze.	Do.
1476	do	13 06 00	145 00 15	1,327	81	83	35.5	co. s. and m. . .	do	
1477	do	13 05 45	145 10 20	1,536	80	83	co. s. and m. . .	do	
1478	do	13 05 30	145 20 30	2,025	80	84	co. s. and m. . .	do	
1479	do	13 05 15	145 30 45	2,675	86	84	35	br. m.	Volcanic mud.	
1480	do	13 05 00	145 40 45	3,071	83	85	br. m. and g. . .	Red clay...	
1481	do	13 04 30	145 49 30	4,472	85	85	br. m. and s. . .	do	
1482	do	12 44 30	145 47 30	5,160	82	85	No specimen	Do.
1483	do	12 40 45	145 56 15	4,249	81	84	br. m. and s. . .	Red clay...	
1484	Nov. 14	13 03 00	145 58 00	4,560	81	84	35.5	No specimen	Do.
1485	do	12 50 15	145 45 00	4,675	82	84	35.6	br. s. and m. . .	Red clay...	Minute spec- imen. Frag- ments of Coccinodis- cus.
1486	do	12 46 00	145 47 30	5,070	81	84	35.9	s.	do	Do.
1487	do	12 44 00	145 46 45	5,101	82	84	36	yl. m.	do	Do.
1488	do	12 43 15	145 49 00	5,269	81	84	No specimen	Deepest sounding.
1489	Nov. 15	13 12 40	145 04 00	1,240	81	84	gy. s. and m. . .	Globigerina ooze.	
1490	do	13 15 45	144 51 30	707	83	84	gy. s. and m. . .	do	
1491	do	13 26 30	145 06 20	939	81	84	gy. s. and m. . .	do	
1492	do	13 35 00	145 13 00	1,054	82	84	36.7	gy. s. and m. . .	do	
1493	do	13 25 45	145 22 30	1,683	84	84	gy. s. and m. . .	do	
1494	do	13 32 30	145 16 00	1,316	82	84	gy. s. and m. . .	do	
1495	do	13 46 15	145 22 45	1,444	80	84	gy. s. and m. . .	do	
1496	do	13 38 30	145 39 45	2,285	81	84	gy. s. gy. and br. m.	Volcanic mud.	Manganese and forami- nifera.
1497	do	13 58 30	145 35 00	1,903	81	84	gy. s. gy. and br. m.	Globigerina ooze.	
1498	Nov. 16	13 56 15	145 57 45	2,259	81	84	gy. and br. m. and s. . . .	Volcanic mud.	
1499	do	14 13 15	145 45 30	2,043	81	84	gy. and br. m. and s. . . .	do	
1500	do	14 17 00	146 05 45	2,650	83	84	br. m. bk. s. . .	Red clay...	
1501	do	14 32 45	145 55 00	2,151	82	84	br. m. bk. s. . .	Volcanic mud.	
1502	do	14 37 30	146 17 30	2,330	81	84	br. m. and g. . .	do	
1503	do	14 53 45	146 07 15	2,253	81	84	g.	do	
1504	Nov. 17	14 58 30	146 29 00	2,586	81	84	br. m. bk. s. . .	Volcanic mud.	No specimen.
1505	do	15 18 00	146 39 45	2,884	81	84	br. m. bk. s. . .	do	
1506	do	15 24 30	146 32 30	2,720	82	84	br. m. bk. s. . .	do	
1507	do	15 30 30	146 38 45	2,983	82	84	35.5	br. m. bk. s. . .	do	
1508	do	15 39 20	146 38 20	2,841	85	84	33	br. m. bk. s. . .	do	
1509	do	15 14 15	146 25 00	2,446	82	84	br. m. bk. s. . .	do	
1510	do	15 32 30	146 47 00	3,167	82	84	34	br. m. bk. s. . .	do	
1511	Nov. 18	15 52 15	146 51 15	2,883	82	84	33	br. m. bk. s. . .	do	
1512	do	15 37 00	147 01 45	2,386	77	84	34	br. m. bk. s. . .	do	
1513	do	15 55 00	147 09 15	2,864	81	84	34	br. m.	Red clay...	
1514	do	15 39 40	147 22 30	2,721	82	84	34	br. m.	do	
1515	do	15 57 30	147 26 30	2,762	82	84	35.2	br. m.	do	
1516	Nov. 19	15 45 00	147 37 00	3,598	82	84	35	br. m.	do	
1517	do	15 45 15	147 41 15	3,996	82	84	35.4	br. m. and s. . .	Volcanic mud.	
1518	do	15 45 45	147 48 30	3,198	81	84	35.5	br. m. and s. . .	do	
1519	do	15 46 20	147 58 15	3,337	83	84	35.4	br. m.	Red clay ..	
1520	do	15 52 00	147 59 00	3,263	83	84	35.6	br. m.	do	
1521	do	15 46 00	148 09 15	2,981	82	84	35.6	br. m. bk. s. . .	Volcanic mud.	
1522	do	16 03 40	147 43 30	2,855	81	84	35.4	R.	Red clay...	Concretions of clay and manganese.
1523	do	16 03 40	147 59 00	2,499	78	84	35.5	R.	do	
1524	Nov. 20	16 03 40	148 14 45	1,587	80	84	36	S.	Globigerina ooze.	
1525	do	15 59 30	148 17 15	1,585	80	84	36.5	S.	do	

Abstract of the official record of soundings—Continued.

GUAM TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
1526	Nov. 20	15 55 30	148 20 15	1,213	81	84	36	S	Globigerina ooze.	
1527	do	15 51 30	148 23 00	2,106	81	84	35.4	S	do	
1528	do	15 47 30	148 26 00	2,391	81	84	35.5	br. m.	Volcanic mud.	
1529	do	15 51 30	148 29 00	1,956	81	84	35	S	Globigerina ooze.	
1530	do	15 38 00	148 04 00	3,096	81	85	35.6	br. m.	Red clay	
1531	do	15 35 20	148 13 00	2,462	81	85	35.2	br. m. bk. s.	Volcanic mud.	
1532	do	15 28 20	148 06 40	2,762	82	84	35	br. m. bk. s.	do	
1533	do	15 26 30	148 17 15	1,731	81	84	35.3	R. G.	do	No specimen.
1534	do	15 29 00	148 27 15	2,280	82	84	35.5	gy. m. and s. bk. s.	Volcanic mud.	
1535	Nov. 21	15 31 60	148 32 45	2,386	81	84	35.5	br. m.	Red clay	
1536	do	15 40 30	148 38 45	1,724	81	84	35.6	R	do	Do.
1537	do	15 50 00	148 44 00	1,081	81	84		wh. s.	Globigerina ooze.	
1538	do	15 42 00	148 50 00	1,710	80	84	35	R. S.	do	Manganese.
1539	do	15 35 30	148 55 45	3,184	83	84	35.5	br. m.	Red clay	
1540	do	15 30 00	148 50 45	3,191	83	84	35.8	br. m.	do	
1541	do	15 19 00	148 20 15	1,747	83	84	35.5	G	Globigerina ooze.	Do.
1542	do	15 10 40	148 05 45	1,397	79	84	35	gy. s.	do	
1543	do	15 06 15	148 14 00	2,942	81	84	35.5	br. m.	Red clay	
1544	do	15 01 40	148 22 15	2,006	79	84	34.8	gy. m. and s.	Globigerina ooze.	
1545	Nov. 22	15 03 45	148 32 30	2,641	76	84	34.6	br. m. bk. s.	Volcanic mud.	No specimen.
1546	do	15 10 30	148 47 30	3,166	78	84	35.4	br. m. bk. s.	do	
1547	do	15 00 00	148 49 45	3,132	78	84	35.1	br. m.	Red clay	
1548	do	14 50 30	148 52 15	3,108	81	84	35.5	br. m.	do	
1549	do	15 00 00	149 04 00	3,135	82	84	35.5	br. m.	do	
1550	do	15 09 00	149 09 30	3,145	80	84	35.4	br. m.	do	
1551	do	14 59 40	149 18 15	3,169	80	84	35.3	br. m.	do	
1552	Nov. 23	14 48 20	149 29 30	3,147	82	84	35.5	br. m.	do	
1553	do	15 07 15	149 33 30	3,170	84	84	35.4	br. m.	do	
1554	do	14 54 30	149 51 30	3,101	83	84	35.3	br. m.	do	
1555	do	15 17 00	149 54 30	3,147	86	85	35.2	br. m.	do	
1556	do	15 05 15	150 07 45	3,214	82	85	35.5	br. m.	do	
1557	do	15 26 45	150 09 10	3,182	81	84		br. m.	do	
1558	Nov. 24	15 15 30	150 24 00	3,206	80	84	35.6	br. m.	do	
1559	do	15 36 20	150 25 30	3,217	82	84	35.5	br. m.	do	
1560	do	15 24 45	150 41 00	3,230	81	84	35.5	br. m.	do	
1561	do	15 44 00	150 47 30	3,240	82	85	35.6	br. m.	do	
1562	do	15 27 00	151 06 30	3,344	82	85	35.4	br. m.	do	
1563	do	15 46 00	151 09 00	3,266	84	84	35.6	br. m.	do	
1564	do	15 30 40	151 24 15	3,177	80	83	35.7	br. m.	do	
1565	Nov. 25	15 49 15	151 28 45	3,289	80	83	35.8	br. m.	do	
1566	do	15 34 45	151 43 30	2,561	83	83	36	br. m.	do	
1567	do	15 38 40	151 45 15	2,672	85	83	35.5	br. m.	do	
1568	do	15 63 30	151 48 30	2,946	82	84	35.5	br. m.	do	
1569	do	16 07 00	151 52 30	825	81	84	39	Co. s. and m.	Globigerina ooze.	
1570	do	16 04 20	151 53 30	815	81	84	39	Co. s. and m.	do	
1571	do	16 01 40	151 54 30	1,348	81	84	36.8	Co. s. and m.	do	
1572	do	15 59 00	151 55 30	1,892	82	84	35.7	Co. s. m. and G.	do	Manganese.
1573	do	15 51 00	151 58 45	2,656	81	84	35.7	br. m.	Red clay	
1574	do	15 51 30	152 04 00	2,757	81	84	35.5	br. m.	do	
1575	do	15 52 15	152 09 15	2,904	81	83	35.5	br. m.	do	
1576	Nov. 26	15 56 00	152 08 30	2,978	80	83	35.7	br. m.	do	
1577	do	16 00 00	152 08 00	2,957	80	83	35.9	br. m.	do	
1578	do	16 03 45	152 07 15	2,778	80	83				No specimen.
1579	do	16 07 30	152 06 45	3,122	80	83	35	br. m.	Red clay	
1580	do	16 11 30	152 05 45	3,121	81	84	35	br. m.	do	
1581	do	16 10 45	152 10 00	3,175	82	84	35	br. m.	do	
1582	do	16 00 00	152 20 15	3,165	86	85	35	br. m.	do	
1583	do	15 44 30	152 04 00	777	85	85	37.5	wh. s. and m.	Globigerina ooze.	
1584	do	16 15 00	152 23 15	3,239	81	84	35	br. m.	Red clay	
1585	do	16 09 00	152 29 15	3,200	80	84	35	br. m.	do	
1586	Nov. 27	16 01 30	152 37 30	3,190	81	83	35	br. m.	do	
1587	do	16 14 30	152 37 00	3,219	80	83	35	br. m.	do	
1588	do	16 24 00	152 34 45	3,288	81	83	34.8	br. m.	do	
1589	do	16 20 30	152 43 45	3,206	82	84	35	br. m.	do	

Abstract of the official record of soundings—Continued.

GUAM TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	<i>Fath- oms.</i>	°	°	°			
1590	Nov. 27	16 26 15	152 51 45	3,195	83	85	35.2	br. m.	Red clay...	
1591	do	16 36 00	152 50 15	3,180	82	84	34.5	br. m.	do	
1592	do	16 32 30	153 00 15	3,197	81	84	35	br. m.	do	
1593	do	16 38 20	153 08 00	3,195	81	84	34.8	br. m.	do	
1594	do	16 48 00	153 06 00	3,194	81	84	35.2	br. m.	do	
1595	Nov. 28	16 44 00	153 16 30	3,193	80	83	35.2	br. m.	do	
1596	do	16 50 15	153 24 00	3,195	79	83	35	br. m.	do	
1597	do	16 58 30	153 22 45	3,191	81	83	35	br. m.	do	
1598	do	16 54 00	153 44 15	2,905	81	84	35	br. m.	do	
1599	do	17 03 30	153 45 00	3,055	79	84	35	br. m.	do	
1600	do	17 13 30	153 47 15	3,110	81	84	35.2	yl. m.	do	Minute speci- men mostly diatoms.
1601	do	17 10 45	153 57 00	1,733	81	84	35	gy. s.	Globigerina ooze.	
1602	do	17 15 45	153 55 15	2,059	81	84	35	yl. m.		No specimen.
1603	do	17 21 30	153 53 30	1,733	80	84	35	gy. s. and m.	Globigerina ooze.	
1604	do	17 22 30	153 48 45	2,923	82	83	35	br. m.	Red clay...	
1605	Nov. 29	17 23 30	153 44 15	3,185	81	83	35.2	br. m.	do	
1606	do	17 29 45	153 45 15	3,185	81	83	35.2	br. m.	do	
1607	do	17 50 30	153 47 15	3,138	81	83	35.2	br. m.	do	
1608	do	17 36 00	154 03 15	3,122	82	84	35	br. m.	do	
1609	do	17 52 00	154 07 15	3,115	82	83	35	br. m.	do	
1610	do	17 36 45	154 21 00	2,977	81	83	35.2	br. m.	do	
1611	do	17 53 40	154 26 00	2,998	81	83	35	br. m.	do	Do.
1612	Nov. 30	17 45 30	154 41 00	3,065	80	83	34.5	br. m.	Red clay...	
1613	do	17 55 00	154 46 00	3,102	80	83	35.2	br. m.	do	
1614	do	17 47 00	154 51 30	3,065	81	83	35.2	br. m.	do	
1615	do	17 39 15	154 57 00	3,112	81	83	35	br. m.	do	
1616	do	17 47 00	155 02 00	3,084	81	83	34	br. m.	do	
1617	do	17 55 45	155 06 45	3,182	81	83	35	br. m.	do	
1618	do	17 47 30	155 12 40	3,138	80	83	35	br. m.	do	
1619	do	17 47 30	155 23 00	3,122	80	83	35	br. m.	do	
1620	do	17 56 00	155 28 00	3,143	78	83	35.1	br. m.	do	
1621	Dec. 1	17 47 15	155 33 30	3,133	80	83	35.5	br. m.	do	
1622	do	17 37 15	155 43 45	3,185	80	83	35.6	br. m.	do	
1623	do	17 57 00	155 48 00	3,164	81	83	35.9	br. m.	do	
1624	do	17 44 30	156 03 15	3,206	79	83	35.2	br. m.	do	
1625	do	18 05 40	156 08 45	3,135	81	83	35.5	br. m.	do	
1626	do	17 54 15	156 25 30	3,193	80	83	35.2	br. m.	do	
1627	Dec. 2	18 15 30	156 30 15	3,159	81	83	35.4	br. m.	do	
1628	do	18 02 45	156 45 15	3,164	81	83	35	br. m.	do	
1629	do	18 23 00	156 52 15	3,188	80	83	35.6	br. m.	do	
1630	do	18 05 00	157 09 15	3,164	80	83	35.2	br. m.	do	
1631	do	18 23 15	157 15 15	3,159	81	83	35.8	br. m.	do	
1632	do	18 08 40	157 29 45	3,168	79	83	35.8	br. m.	do	
1633	Dec. 3	18 26 40	157 36 00	3,173	80	82	35.2	br. m.	do	
1634	do	18 12 00	157 51 00	3,190	80	82	35	br. m.	do	
1635	do	18 32 30	157 54 45	3,164	82	83	35.5	br. m.	do	Do.
1636	do	18 20 45	158 11 45	3,185	82	83	35.5	br. m.	Red clay...	
1637	do	18 39 20	158 16 15	3,159	78	82	35	br. m.	do	Do.
1638	Dec. 4	18 25 30	158 31 30	3,163	80	83	35	br. m.	Red clay...	
1639	do	18 46 00	158 37 30	3,106	79	83	35.2	br. m.	do	
1640	do	18 31 00	158 53 15	3,170	81	83	35.2	br. m.	do	
1641	do	18 50 00	158 58 20	3,158	80	83	35	br. m.	do	
1642	do	18 54 00	159 14 30	3,130	80	83	34.5	br. m.	do	
1643	do	18 52 30	159 20 00	3,148	79	82	35.3	br. m.	do	
1644	do	18 38 00	159 35 30	3,218	79	82	35	br. m.	do	
1645	Dec. 5	18 56 30	159 40 00	3,164	79	82	35	br. m.	do	
1646	do	18 42 00	159 56 45	3,131	80	82	35	br. m.	do	
1647	do	19 00 30	160 01 30	3,150	80	82	35	br. m.	do	
1648	do	18 51 15	160 15 30	3,126	80	82	35	br. m.	do	
1649	do	19 16 40	160 14 45	2,769	79	82	35	br. m.	do	
1650	Dec. 6	19 07 00	160 30 45	3,038	80	82	35	br. m.	do	
1651	do	19 22 00	160 34 30	2,567	79	82	35	br. m.	do	
1652	do	19 23 00	160 44 30	2,886	79	82	35	br. m.	do	
1653	do	19 32 00	160 40 15	2,584	80	82	35	br. m.	do	
1654	do	19 40 30	160 44 45	2,673	80	82	35	br. m.	do	
1655	do	19 40 30	160 55 45	2,912	80	83	35	br. m.	do	
1656	do	19 48 00	160 51 30	2,859	81	83	35	br. m.	do	
1657	do	19 54 45	160 58 30	2,854	80	83	35	br. m.	do	
1658	do	19 54 00	161 09 15	2,912	79	83	35	br. m.	do	
1659	do	20 04 00	161 06 30	2,468	79	82	35	br. m.	do	Do.
1660	Dec. 7	20 11 45	161 13 00	2,270	79	82	35	br. m.	do	Do.
1661	do	20 10 30	161 24 15	2,751	79	82	35	br. m.	Red clay...	

Abstract of the official record of soundings—Continued.

GUAM TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
1662	Dec. 7	20 20 30	161 21 00	2,596	78	82	br. m. and s.	Red clay...	
1663	do	20 30 30	161 18 00	2,552	79	82	br. m.	do	
1664	do	20 29 00	161 28 45	2,788	81	82	34.2	br. m.	do	
1665	do	20 27 00	161 39 45	2,827	79	82	35	br. m.	do	
1666	do	20 34 53	161 37 30	2,848	79	82	br. m.	do	
1667	do	20 40 00	161 46 30	2,817	79	82	36	br. m.	do	
1668	do	20 35 30	161 56 00	2,807	79	82	35	br. m.	do	
1669	do	20 45 20	161 55 45	2,816	79	82	br. m.	do	
1670	Dec. 8	20 51 00	162 05 00	2,838	79	82	35	br. m.	do	
1671	do	20 46 20	162 14 30	2,817	78	81	br. m.	do	No specimen.
1672	do	20 56 40	162 14 30	2,819	78	81	br. m.	Red clay	
1673	do	21 08 00	162 14 30	2,815	80	82	br. m.	do	
1674	do	20 58 30	162 33 30	2,832	79	82	br. m.	do	
1675	do	21 14 45	162 41 30	2,114	80	82	35	br. m. and s.	do	
1676	do	21 11 40	162 45 15	1,087	79	82	wh. m. gy. s.	Globigerina ooze.	
1677	do	21 10 30	162 47 15	1,054	79	82	37.3	gy. m. and s.	do	
1678	do	21 08 00	162 49 45	1,283	78	82	38	gy. m. and s.	do	
1679	do	21 03 45	162 53 06	2,203	78	82	br. m. and s.	do	Do.
1680	do	20 59 00	162 56 15	2,451	78	82	35	br. m.	Red clay	
1681	do	20 54 40	162 59 45	2,498	79	81	br. m.	do	
1682	Dec. 9	20 58 20	163 03 15	2,270	78	81	35.1	br. m.	do	
1683	do	21 02 00	163 06 45	1,798	78	81	35.2	gy. m.	Globigerina ooze.	
1684	do	21 05 45	163 10 30	828	78	81	37	gy. s.	do	
1685	do	21 01 40	163 14 15	1,507	78	81	gy. s.	do	
1686	do	20 57 30	163 17 45	2,003	78	81	35.2	gy. s.	do	
1687	do	20 53 00	163 20 30	2,435	78	81	br. m.	Red clay	
1688	do	20 48 00	163 21 15	2,630	80	82	35	br. m.	do	
1689	do	20 55 30	163 28 45	2,556	83	82	br. m.	do	
1690	do	21 01 00	163 27 45	1,992	79	82	br. m. and s.	Globigerina ooze.	
1691	do	21 06 15	163 26 45	1,912	79	82	35	gy. m. and s.	do	
1692	do	21 14 40	163 18 15	1,754	79	82	gy. m. and s.	do	
1693	do	21 23 00	163 10 00	2,050	77	82	35	gy. s.	do	
1694	do	21 26 40	163 06 00	2,272	77	81	br. m.	Red clay	
1695	do	21 29 45	163 02 45	2,377	79	81	35	br. m.	do	
1696	do	21 33 00	162 59 00	2,050	78	81	35	gy. m.	Globigerina ooze.	
1697	Dec. 10	21 36 20	162 55 30	2,103	79	81	35	wh. and bk. s.	do	
1698	do	21 30 40	162 55 15	2,093	78	81	35.2	gy. m.	do	
1699	do	21 28 30	162 55 00	1,456	77	80	gy. m. and s.	do	
1700	do	21 20 00	162 54 45	1,917	78	80	gy. m. and s.	do	
1701	do	21 22 40	162 50 40	1,352	76	80	gy. m. and s.	do	
1702	do	21 24 45	162 46 40	1,854	75	81	gy. m. and s.	do	
1703	do	21 27 00	162 43 00	2,731	76	81	br. m.	Red clay	
1704	do	21 21 00	162 42 00	1,879	79	81	gy. m. and s.	Globigerina and G. ooze.	Largemanganesenodule.
1705	do	21 15 15	162 41 00	1,083	78	81	gy. m. and s.	do	
1706	do	21 13 00	162 44 40	1,061	79	81	36	gy. m.	do	
1707	do	21 16 00	162 38 00	2,029	79	81	35	gy. m.	do	
1708	do	21 19 15	162 33 30	2,953	79	82	br. m.	Red clay	No specimen.
1709	do	21 13 30	162 27 45	2,900	78	82	br. m.	Red clay	
1710	do	21 07 40	162 22 20	2,867	78	81	br. m.	Diatom ooze.	Coscinodiscus rex.
1711	do	21 16 45	162 17 45	2,827	78	81	br. m.	Red clay	
1712	Dec. 11	21 24 15	162 24 00	2,879	78	81	br. m.	Red clay	No specimen.
1713	do	21 27 00	162 34 00	2,879	77	80	br. m.	do	
1714	do	21 37 00	162 33 00	2,808	77	80	br. m.	do	
1715	do	21 35 15	162 43 00	2,912	79	81	br. m.	do	
1716	do	21 38 20	162 50 00	2,881	78	81	br. m.	do	
1717	do	21 47 00	162 45 20	2,570	78	81	br. m.	do	
1718	do	21 43 30	162 49 45	2,854	78	81	br. m.	do	
1719	do	21 41 40	162 57 00	2,830	78	81	br. m.	do	
1720	do	21 46 30	162 57 00	2,798	78	81	br. m.	do	
1721	do	21 56 45	162 57 00	2,890	77	80	br. m.	do	
1722	do	21 48 30	163 01 15	2,831	76	80	br. m.	do	
1723	Dec. 12	21 44 00	163 03 15	2,774	77	81	br. m.	do	
1724	do	21 39 15	163 05 30	2,676	77	81	gy. m.	Diatom ooze.	Coscinodiscus rex.
1725	do	21 43 30	163 07 00	2,628	77	81	br. m.	Red clay	
1726	do	21 48 00	163 09 00	2,120	78	81	br. m.	do	
1727	do	21 52 00	163 10 45	2,360	79	81	br. m.	do	
1728	do	21 56 30	163 11 30	2,868	79	81	br. m.	do	
1729	do	22 01 45	163 11 30	2,985	79	81	br. m.	do	

Abstract of the official record of soundings—Continued.

GUAM TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1899.	° ' "	° ' "	Fath- oms.	°	°	°			
1730	Dec. 12	21 53 30	163 16 15	1,900	79	81	gy. m.	Globigernia ooze.	
1731do....	21 58 00	163 20 45	2,649	81	81	br. m.	Red clay...	
1732do....	21 46 45	163 21 15	2,241	79	81	br. m.do....	
1733do....	21 40 20	163 20 30	2,672	79	81	br. m.do....	
1734do....	21 35 00	163 20 00	2,738	78	81	br. m.do....	
1735do....	21 29 20	163 19 15	2,612	78	80	br. m.do....	
1736do....	21 33 20	163 24 00	2,775	78	80	br. m.do....	
1737	Dec. 13	21 37 20	163 28 40	2,876	78	80	br. m.do....	
1738do....	21 41 00	163 33 20	2,943	78	80	br. m.do....	
1739do....	21 45 20	163 37 45	2,966	78	80	br. m.do....	
1740do....	21 39 20	163 38 45	2,950	80	81	br. m.do....	
1741do....	21 34 00	163 39 45	2,945	80	81	br. m.do....	
1742do....	21 33 40	163 13 00	2,395	80	81	br. m.do....	
1743do....	21 39 30	163 11 30	2,289	79	81	br. m.do....	
1744do....	21 45 30	163 10 30	2,299	79	81	br. m.do....	
1745do....	21 47 30	163 14 20	1,696	79	81	gy. m. and s.	Globigerina ooze.	
1746do....	21 43 30	163 13 45	1,973	78	81	bk. and wh. s. and bk. G.do....	Manganese.
1747do....	21 39 15	163 13 20	2,225	78	81	br. m.	Red clay...	
1748	Dec. 14	21 41 30	163 56 15	3,033	78	80	br. m.do....	
1749do....	21 47 00	164 05 15	2,967	77	80	br. m.do....	
1750do....	21 42 00	164 10 15	2,974	78	81	br. m.do....	
1751do....	21 31 00	164 26 30	3,021	78	81	br. m.do....	
1752do....	21 51 30	163 58 30	3,000	78	82	br. m.do....	
1753do....	21 56 15	164 16 45	2,902	77	82	br. m.do....	
1754do....	21 44 30	164 34 45	3,029	77	80	br. m.do....	
1755	Dec. 15	22 05 15	164 39 45	3,036	77	80	br. m.do....	
1756do....	21 53 15	164 57 15	3,018	76	80	br. m.do....	
1757do....	22 13 45	165 02 00	3,085	80	81	br. m.do....	
1758do....	22 01 30	165 22 20	3,078	82	82	br. m.do....	
1759do....	22 20 00	165 20 20	3,107	78	82	br. m.do....	
1760do....	22 05 30	165 33 45	3,070	77	82	br. m.do....	
1761do....	22 25 30	165 35 15	3,234	76	81	br. m.do....	
1762	Dec. 16	22 10 45	165 47 15	3,198	76	80	br. m.do....	
1763do....	22 30 45	165 49 30	3,229	78	81	br. m.do....	
1764do....	22 12 20	166 88 30	3,227	78	81	br. m.do....	
1765do....	22 28 20	166 10 30	3,126	78	82	br. m.do....	No specimen.
1766do....	22 13 30	166 26 30	3,237	79	82	br. m.do....	
1767do....	22 32 40	166 31 45	3,269	79	81	br. m.do....	
1768	Dec. 17	22 17 00	166 47 00	3,235	79	80	br. m.do....	
1769do....	22 36 40	166 52 45	3,228	75	80	br. m.do....	
1770do....	22 20 20	167 10 00	3,206	77	80	br. m.do....	
1771do....	22 44 40	167 13 00	3,261	74	80	br. m.do....	
1772do....	22 32 45	167 21 45	3,208	75	80	br. m.do....	
1773do....	22 50 30	167 26 00	3,321	73	79	br. m.do....	
1774	Dec. 18	22 38 30	167 41 45	3,164	74	79	br. m.do....	
1775do....	22 59 00	167 48 30	3,310	71	78	br. m.do....	Do.
1776do....	22 46 30	168 02 00	3,379	73	79	br. m.	Red clay ..	
1777do....	22 59 00	168 03 00	3,261	73	80	br. m.do....	
1778do....	22 46 15	168 15 45	3,298	73	79	br. m.do....	
1779do....	23 05 00	168 16 45	3,207	72	78	br. m.do....	
1780do....	22 53 20	168 34 00	3,169	71	78	br. m.do....	
1781	Dec. 19	23 12 15	168 35 40	3,251	72	78	br. m.do....	
1782do....	23 00 15	168 52 45	3,119	71	77	br. m.do....	
1783do....	23 18 30	168 54 15	3,221	73	80	br. m.do....	
1784do....	23 09 00	169 07 15	3,256	73	80	br. m.do....	
1785do....	23 28 40	169 12 30	3,288	73	80	br. m.do....	
1786do....	23 18 30	169 30 15	3,238	72	79	br. m.do....	Do.
1787do....	23 37 40	169 34 30	3,320	72	78	br. m.	Red clay ..	
1788	Dec. 20	23 28 00	169 54 30	3,318	73	77	br. m.do....	
1789do....	23 47 30	169 56 00	3,331	75	78	br. m.do....	
1790do....	23 38 00	170 11 45	3,288	77	79	br. m.do....	
1791do....	24 05 00	170 07 20	3,243	76	79	br. m.do....	
1792do....	23 49 30	170 29 20	3,247	75	79	br. m.do....	
1793do....	24 05 40	170 35 00	3,273	73	78	br. m.do....	Do.
1794	Dec. 21	23 54 20	170 56 00	3,250	72	77	br. m.do....	Do.
1795do....	24 09 45	171 00 45	3,257	69	76	br. m.	Red clay ..	
1796do....	24 15 15	170 56 30	3,257	69	76	br. m.do....	
1797do....	24 11 00	171 11 15	3,265	70	76	br. m.do....	
1798do....	24 34 00	171 06 30	3,252	69	76	br. m.do....	
1799do....	24 21 00	171 29 15	3,196	70	76	br. m.do....	
1800	Dec. 22	24 39 45	171 30 00	3,214	68	75	br. m.do....	
1801do....	24 30 30	171 51 40	3,206	68	75	br. m.do....	
1802do....	24 49 30	171 49 40	3,281	68	75	br. m.do....	
1803do....	24 37 30	172 08 40	3,198	69	76	br. m.do....	

Abstract of the official record of soundings—Continued.

GUAM TO MIDWAY ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude east.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1899.	° ' "	° ' "	Fathoms.	°	°	°			
1804	Dec. 22	24 55 20	172 11 00	3,225	67	75	br. m.	Red clay...	
1805do....	24 45 20	172 29 00	3,258	68	75	br. m.do....	
1806do....	25 07 00	172 32 20	3,432	69	75	br. m.do....	
1807	Dec. 23	24 55 30	172 50 00	3,237	68	75	br. m.do....	
1808do....	24 47 15	172 53 30	3,202	68	76	br. m.do....	
1809do....	25 08 40	172 56 30	3,208	68	76	br. m.do....	
1810do....	24 56 45	173 13 15	3,156	69	75	br. m.do....	
1811do....	25 17 00	173 17 45	3,227	69	75	br. m.do....	
1812	Dec. 24	25 05 00	173 34 30	3,206	68	75	br. m.do....	
1813do....	25 25 30	173 38 45	3,231	68	75	br. m.do....	
1814do....	25 13 30	173 55 00	3,258	70	76	br. m.do....	
1815do....	25 26 40	173 55 30	3,283	70	75	br. m.do....	
1816do....	25 14 30	174 13 00	3,257	70	75	br. m.do....	
1817do....	25 35 00	174 20 00	3,257	67	74	br. m.do....	
1818	Dec. 25	25 23 00	174 36 30	3,208	69	75	br. m.do....	
1819do....	25 44 15	174 43 30	3,475	70	75	br. m.do....	
1820do....	25 26 00	175 01 15	3,258	72	75	br. m.do....	
1821	Dec. 26	25 47 30	175 12 00	3,434	78	75	br. m.do....	
1822	Dec. 28	26 01 00	175 47 00	3,252	68	74	br. m.do....	
1823do....	25 54 40	175 30 00	3,227	69	75	br. m.do....	
1824do....	25 36 00	175 23 45	3,276	68	75	br. m.do....	
1825do....	25 39 30	175 43 15	3,292	70	75	br. m.do....	
1826	Dec. 29	25 46 15	176 05 00	3,083	69	75	br. m.do....	
1827do....	26 06 40	176 09 20	3,357	67	73	br. m.do....	
1828do....	25 54 00	176 26 30	3,230	69	73	br. m.do....	
1829do....	26 14 00	176 30 00	3,242	70	75	br. m.do....	
1830do....	26 02 30	176 46 20	3,194	70	74	br. m.do....	
1831	Dec. 30	26 24 00	176 49 30	3,119	70	75	br. m.do....	
1832do....	26 12 45	177 06 00	3,086	70	75	br. m.do....	
1833do....	26 34 00	177 09 00	3,074	72	75	br. m.do....	
1834do....	26 17 00	177 32 15	3,229	70	75	br. m.do....	
1835do....	26 38 20	177 31 15	3,061	67	75	br. m.do....	
1836do....	26 18 30	177 48 15	3,013	67	75	br. m.do....	
1837	Dec. 31	26 37 40	177 52 45	3,115	63	75	br. m.do....	
1838do....	26 24 00	178 09 15	3,105	67	74	br. m.do....	
1839do....	26 43 30	178 14 30	3,047	63	74	br. m.do....	
1840do....	26 29 45	178 33 30	2,876	72	74	br. m.do....	
1841do....	26 49 30	178 39 00	3,007	63	72	br. m.do....	
1842do....	26 36 15	178 56 30	3,078	64	73	br. m.do....	
	1900.									
1843	Jan. 1	26 55 45	179 02 45	3,022	65	72	br. m.do....	
1844do....	26 43 15	179 20 30	3,038	66	72	br. m.do....	
1845do....	27 02 45	179 27 00	2,951	70	73	br. m.do....	
1846do....	26 47 45	179 43 45	2,970	68	71	br. m.do....	
1847do....	27 06 40	179 50 30	2,993	68	71	br. m.do....	
			West.							
1848do....	26 52 00	179 55 30	2,947	66	71	br. m.do....	
1849do....	27 10 30	179 46 00	2,939	65	70	br. m.do....	
1850do....	26 55 40	179 34 00	3,036	65	70	br. m.do....	
1851do....	27 13 15	179 22 30	2,951	64	70	br. m.do....	
1852do....	26 58 30	179 12 30	2,951	65	71	br. m.do....	
1853do....	27 16 45	179 06 00	2,915	64	70	br. m.do....	
1854do....	27 02 00	178 51 30	2,997	67	70	br. m.do....	
1855do....	27 21 15	178 46 30	2,895	64	68	br. m.do....	
1856	Jan. 2	27 06 30	178 30 15	2,859	69	68	br. m.do....	
1857do....	27 26 45	178 25 30	2,772	65	67	br. m.do....	
1858do....	27 12 00	178 10 00	2,757	69	70	br. m.do....	
1859do....	27 30 00	178 04 15	2,437	67	70	br. m.do....	
1860do....	27 11 30	177 56 00	2,734	67	70	br. m.do....	
1861do....	27 21 00	177 37 00	2,737	67	70	br. m.do....	
1862	Jan. 3	27 33 00	177 48 30	2,462	66	70	br. m.do....	
1863do....	27 44 15	177 25 30	2,470	67	70	br. m.do....	
1864do....	27 53 00	177 25 45	2,224	67	69	gy. m.	Globigerina ooze.	
1865do....	27 57 00	177 15 30	2,185	69	69	gy. m.do....	No specimen.
1866do....	28 07 00	177 22 15	1,503	66	70	g.	Globigerina ooze.	Manganese.
1867do....	27 57 00	177 34 00	2,311	68	68	gy. m.do....	
1868do....	28 05 00	177 32 00	1,624	67	66	gy. m. and s.do....	
1869do....	28 10 00	177 26 30	57	67	66do....	No specimen.
1870do....	28 10 30	177 31 15	1,618	66	66	gy. m. and s.	Globigerina ooze.	
1871do....	28 17 30	177 28 15	325	67	66do....	Nospecimen, Midway Islands.

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO HAWAIIAN ISLANDS.

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
1872	1900. Jan. 3	° ' "	° ' "	Fath- oms.	°	°	°	gy. m. and s.	Globigerina ooze.	
1873	do	28 27 20	177 31 45	1,767	66	66	66	gm. m. and s.	do	
1874	do	28 27 20	177 13 30	2,188	66	66	66	gm. m. and s.	do	
1875	Jan. 4	28 45 15	177 29 45	2,884	64	65	65	br. m.	Red clay	
1876	do	28 43 30	177 15 30	2,887	63	66	66	br. m.	do	
1877	do	28 51 20	177 07 30	2,941	62	66	66	br. m.	do	
1878	do	29 00 20	177 14 00	3,043	62	66	66	br. m.	do	
1879	do	28 57 20	176 59 30	2,973	63	69	69	br. m.	do	
1880	do	29 00 45	176 49 00	2,943	63	69	69	br. m.	do	
1881	do	29 08 45	176 54 15	3,002	64	68	68	br. m.	do	
1882	do	29 03 00	176 38 00	2,951	61	68	68	br. m.	do	
1883	Jan. 5	29 11 00	176 32 30	2,978	62	67	67	br. m.	do	
1884	do	29 00 30	176 27 30	2,936	63	69	69	br. m.	do	
1885	do	28 51 20	176 25 45	2,797	62	69	69	br. m.	do	
1886	do	28 57 30	176 17 15	2,871	66	68	68	br. m.	do	
1887	do	29 05 30	176 11 30	2,891	66	70	70	br. m.	do	
1888	do	28 54 20	176 07 00	2,787	66	69	69	br. m.	do	
1889	do	28 55 00	175 54 00	2,818	67	70	70	br. m.	do	
1890	do	28 46 30	175 55 15	2,754	67	69	69	br. m.	do	
1891	do	28 32 15	175 51 00	2,655	69	69	69	br. m.	do	
1892	Jan. 8	28 49 40	175 32 15	2,797	64	69	69	br. m.	do	Jan. 6 and 7, "Riding out gale."
1893	do	28 26 45	175 28 30	2,576	67	69	69	br. m.	do	
1894	do	28 41 15	175 13 45	2,860	68	69	69	br. m.	do	
1895	do	28 21 00	175 09 00	2,838	67	69	69	br. m.	do	
1896	Jan. 9	28 35 20	174 54 00	2,952	69	69	69	br. m.	do	
1897	do	28 15 15	174 49 00	2,931	68	69	69	br. m.	do	
1898	do	28 30 00	174 34 15	2,951	69	69	69	br. m.	do	
1899	do	28 11 20	174 27 30	3,035	71	68	68	br. m.	do	
1900	do	28 24 30	174 12 30	2,956	69	68	68	br. m.	do	
1901	do	28 02 45	174 08 15	2,952	68	67	67	br. m.	do	
1902	Jan. 10	28 12 45	173 48 00	2,983	68	67	67	br. m.	do	
1903	do	27 51 00	173 44 00	3,020	65	67	67	br. m.	do	
1904	do	28 01 15	173 24 15	2,887	64	67	67	br. m.	do	
1905	do	27 46 30	173 22 45	2,914	67	66	66	br. m.	do	
1906	do	27 59 20	173 03 20	2,810	63	66	66	br. m.	do	
1907	do	27 40 30	173 01 00	2,797	63	66	66	br. m.	do	
1908	Jan. 11	27 54 15	172 41 45	2,774	64	65	65	br. m.	do	
1909	do	27 35 30	172 36 45	2,764	65	66	66	br. m.	do	
1910	do	27 47 00	172 22 45	2,746	66	68	68	br. m.	do	
1911	do	27 26 00	172 18 30	2,746	67	71	71	br. m.	do	
1912	do	27 39 00	172 02 45	2,732	67	70	70	br. m.	do	
1913	do	27 18 30	171 58 15	2,727	69	70	70	br. m.	do	
1914	Jan. 12	27 31 45	171 41 15	2,710	70	70	70	br. m.	do	
1915	do	27 11 40	171 37 00	2,710	71	70	70	br. m.	do	
1916	do	27 25 00	171 21 30	2,689	71	71	71	br. m.	do	
1917	do	27 05 30	171 16 30	2,731	73	71	71	br. m.	do	
1918	do	27 18 45	170 58 15	2,603	74	72	72	br. m.	do	
1919	do	27 08 20	170 56 00	2,607	71	71	71	br. m.	do	
1920	do	27 15 15	170 48 15	2,581	71	71	71	br. m.	do	
1921	do	27 09 00	170 48 45	2,593	71	71	71	br. m.	do	
1922	do	27 03 00	170 49 00	2,265	71	70	70	br. m.	do	
1923	Jan. 13	27 07 00	170 45 15	2,598	72	70	70	br. m.	do	
1924	do	27 11 00	170 42 00	2,573	71	70	70	br. m.	do	
1925	do	26 49 20	170 41 15	2,607	72	71	71	br. m.	do	
1926	do	27 01 00	170 26 00	2,535	78	72	72	br. m.	do	
1927	do	26 43 20	170 21 00	2,597	79	72	72	br. m.	do	
1928	do	26 57 00	170 02 45	2,564	75	73	73	br. m.	do	
1929	do	26 39 00	169 54 45	2,500	72	73	73	br. m.	do	
1930	Jan. 14	26 54 30	169 36 00	2,460	71	72	72	br. m.	do	
1931	do	26 36 00	169 29 00	2,528	71	72	72	br. m.	do	
1932	do	26 52 00	169 10 00	2,458	72	72	72	br. m.	do	
1933	do	26 40 30	169 06 45	2,504	73	72	72	br. m.	do	
1934	do	26 43 00	168 55 45	2,499	74	73	73	br. m.	do	
1935	do	26 47 00	168 48 30	2,458	74	73	73	br. m.	do	
1936	do	26 36 30	168 46 45	2,501	72	73	73	br. m.	do	
1937	do	26 34 00	168 36 15	2,507	71	72	72	br. m.	do	
1938	do	26 40 30	168 27 45	2,434	73	72	72	br. m.	do	
1939	do	26 30 45	168 26 15	2,481	68	72	72	br. m.	do	
1940	Jan. 15	26 36 30	168 17 30	2,499	64	71	71	br. m.	Red clay	No specimen.
1941	do	26 31 20	168 08 30	2,787	62	70	70	br. m.	do	
1942	Jan. 16	26 22 45	168 08 15	2,562	62	70	70	br. m.	do	
1943	do	26 17 00	167 58 15	2,466	62	70	70	br. m.	do	
1944	do	26 23 00	167 49 45	2,540	62	69	69	br. m. bk. s.	do	
1945	do	26 13 45	167 48 30	2,529	62	70	70	br. m.	do	

Abstract of official record of soundings—Continued.

MIDWAY ISLANDS TO HAWAIIAN ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Surface.	Bottom.			
	1900.	° ' "	° ' "	Fathoms.	°	°	°			
1946	Jan. 16	26 09 00	167 39 45	2,555	62	70	br. m.	Red clay...	
1947	...do...	26 14 20	167 29 45	2,550	62	70	br. m.do...	
1948	Jan. 17	26 05 30	167 29 30	2,620	62	69	br. m.do...	
1949	...do...	26 00 45	167 20 30	2,666	65	70	br. m. bk. s.do...	
1950	...do...	26 05 45	167 10 30	2,694	71	71	br. m.do...	
1951	...do...	25 56 45	167 10 45	2,693	76	71	br. m. br. and bk. s.do...	
1952	...do...	25 52 00	167 11 00	2,682	71	71	G.do...	Manganese.
1953	...do...	25 47 00	167 10 45	2,252	68	71	br. m.do...	
1954	...do...	25 50 00	167 06 30	2,719	66	71	br. m. br. and bk. s.do...	
1955	...do...	25 52 45	167 02 00	2,706	67	71	br. m.do...	
1956	...do...	25 55 30	166 57 15	2,708	64	70	br. m.do...	
1957	...do...	25 46 00	167 00 45	2,728	66	70	br. m.do...	
1958	...do...	25 41 15	167 02 30	2,751	67	70	gy. s. and m.do...	No specimen.
1959	...do...	25 44 15	166 53 30	2,716	70	70	br. m.do...	Do.
1960	Jan. 18	25 47 15	166 44 45	2,731	69	71	br. m.	Red clay...	
1961	...do...	25 37 20	166 43 45	2,788	71	72	br. m.do...	
1962	...do...	25 24 40	166 40 30	2,807	73	72	br. m.do...	
1963	...do...	25 35 45	166 27 15	2,702	69	71	br. m.do...	
1964	...do...	25 16 20	166 22 00	2,769	69	71	br. m.do...	
1965	...do...	25 30 45	166 06 15	2,748	69	71	br. m.do...	
1966	...do...	25 12 30	166 01 00	2,799	69	72	br. m.do...	
1967	Jan. 19	25 25 00	165 45 00	2,695	72	73	br. m.do...	
1968	...do...	25 06 20	165 40 45	2,782	76	73	br. m.do...	
1969	...do...	25 19 20	165 23 15	2,705	71	72	br. m.do...	
1970	...do...	24 59 30	165 20 45	2,758	70	71	br. m.do...	
1971	...do...	25 12 15	165 05 15	2,722	69	71	br. m.do...	
1972	...do...	24 53 20	165 02 30	2,760	70	72	br. m.do...	
1973	Jan. 20	25 06 15	164 46 00	2,745	74	73	br. m.do...	
1974	...do...	24 47 40	164 44 00	2,874	71	75	br. m.do...	
1975	...do...	25 00 00	164 28 00	2,744	70	74	br. m.do...	
1976	...do...	24 37 00	164 24 20	2,745	71	75	br. m.do...	
1977	...do...	24 50 40	164 12 30	2,721	70	74	br. m.do...	
1978	...do...	24 30 40	164 07 15	2,711	71	74	br. m.do...	
1979	Jan. 21	24 45 00	163 54 45	2,725	71	73	br. m.do...	
1980	...do...	24 27 00	163 49 15	2,725	70	73	br. m.do...	
1981	...do...	24 43 00	163 36 00	2,741	71	74	br. m.do...	
1982	...do...	24 21 00	163 28 00	2,769	75	74	br. m.do...	
1983	...do...	24 35 00	163 13 45	2,741	72	74	br. m.do...	
1984	...do...	24 16 00	163 06 45	2,718	74	73	br. m.do...	
1985	Jan. 22	24 30 00	162 51 30	2,746	73	73	br. m.do...	
1986	...do...	24 10 40	162 46 45	2,705	73	74	br. m.do...	
1987	...do...	24 23 00	162 30 00	2,710	78	74	br. m.do...	
1988	...do...	24 05 30	162 28 15	2,638	77	74	br. m.do...	
1989	...do...	24 16 20	162 06 45	2,626	74	75	br. m.do...	
1990	...do...	23 58 00	162 04 15	2,473	71	74	br. m.do...	
1991	...do...	24 10 00	161 43 45	2,545	70	74	br. m.do...	
1992	Jan. 23	23 50 30	161 40 00	2,432	70	74	br. m.	Red clay...	
1993	...do...	24 03 30	161 20 15	2,596	70	74	br. m.do...	
1994	...do...	23 43 00	161 16 15	2,492	73	75	br. m.do...	
1995	...do...	23 54 45	161 01 30	2,607	74	76	br. m.do...	
1996	...do...	23 35 00	161 00 30	2,605	75	75	br. m.do...	
1997	...do...	23 47 20	160 45 30	2,638	70	74	br. m.do...	
1998	...do...	23 27 30	160 45 00	2,656	71	74	br. m.do...	
1999	Jan. 24	23 40 30	160 28 45	2,638	69	74	br. m.do...	
2000	...do...	23 20 40	160 28 00	2,638	72	75	br. m.do...	
2001	...do...	23 33 30	160 09 15	2,645	76	75	br. m.do...	
2002	...do...	23 13 30	160 08 15	2,679	74	76	br. m.do...	
2003	...do...	23 21 15	159 49 45	2,712	70	75	br. m.do...	
2004	...do...	23 01 45	159 52 00	2,702	71	75	br. m.do...	
2005	Jan. 25	23 08 45	159 30 15	2,689	71	75	br. m.do...	
2006	...do...	22 50 20	159 34 30	2,411	70	75	br. m.do...	
2007	...do...	22 53 40	159 21 15	2,586	68	74	br. m.do...	
2008	...do...	22 57 20	159 09 30	2,659	69	74	br. m.do...	
2009	...do...	23 01 30	159 06 15	2,659	69	75	br. m.do...	No specimen
2010	...do...	22 52 15	159 08 00	2,429	68	75	br. m.	Red clay...	
2011	...do...	22 48 00	159 07 00	2,468	69	75	br. m.do...	
2012	...do...	22 48 00	159 07 00	2,400	68	75	br. m.do...	
2013	...do...	22 48 00	159 00 15	2,535	69	74	br. m.do...	
2014	...do...	22 55 20	158 49 15	2,633	66	74	br. m.do...	
2015	Jan. 26	22 46 40	158 50 00	2,556	67	74	br. m.do...	
2016	...do...	22 42 40	158 41 00	2,670	66	75	br. m.do...	
2017	...do...	22 50 00	158 32 40	2,638	66	75	br. m.do...	

Abstract of the official record of soundings—Continued.

MIDWAY ISLANDS TO HAWAIIAN ISLANDS—Continued.

Station No.	Date.	Latitude north.	Longitude west.	Depth.	Temperatures.			Character of bottom.	Deposit.	Remarks.
					Air.	Sur- face.	Bot- tom.			
	1900.	° ' "	° ' "	Fath- oms.	°	°	°			
2018	Jan. 26	22 42 00	158 34 00	2,658	66	75	br. m.	Red clay...	
2019	...do...	22 37 45	158 30 00	2,676	66	76	br. m.do....	
2020	...do...	22 34 00	158 32 15	2,726	67	76	br. m.do....	
2021	...do...	22 26 40	158 33 40	2,710	69	76	br. m.do....	
2022	...do...	22 33 30	158 47 00	2,329	69	76	br. m.do....	
2023	...do...	22 24 00	158 42 20	2,763	68	74	br. m.do....	
2024	...do...	22 21 00	158 39 00	2,570	67	74	br. m.do....	
2025	...do...	22 21 45	158 32 30	2,670	68	74	br. m.do....	
2026	Jan. 27	22 29 45	158 23 00	2,705	69	74	br. m.do....	
2027	...do...	22 22 00	158 25 30	2,715	69	74	br. m.do....	
2028	...do...	22 19 30	158 18 30	2,370	69	74	br. m.do....	
2029	...do...	22 11 00	158 17 30	2,518	70	74	br. m.do....	
2030	...do...	22 08 30	158 16 20	2,519	70	74	br. m.do....	
2031	...do...	21 54 30	158 11 45	1,624	72	74	gy. m. and s.	Globigerina ooze.	
2032	...do...	21 48 20	158 10 45	1,014	71	76	gy. m. and s.	...do....	
2033	...do...	21 43 45	158 09 00	249	70	76	gy. m. and s.	...do....	
2034	...do...	21 41 45	158 08 40	175	70	76	gy. m. and s.	...do....	
2035	...do...	21 40 42	158 07 27	114	gy. m. and s.	...do....	
2036	...do...	21 40 32	158 07 20	82	gy. m. and s.	Coral sand.	
2037	...do...	21 39 46	158 06 25	55	gy. and bk. s.	...do....	
2038	...do...	21 39 20	158 05 55	34	gy. and bk. s.	...do....	
2039	...do...	21 39 08	158 05 37	24	gy. and br. s.	...do....	Long threads of volcanic glass, Wai- mea Bay.
2040	...do...	21 39 03	158 05 28	21	gy. and br. s.	...do....	
2041	...do...	21 40 00	158 05 35	33	gy. and br. s.	...do....	
2042	...do...	21 41 00	158 05 46	55	gy. and br. s.	...do....	
2043	...do...	21 41 30	158 05 50	58	gy. and br. s.	...do....	
2044	...do...	21 42 00	158 05 55	69do....	No specimen.
2045	...do...	21 42 35	158 06 00	93	gy. and br. s.	Coral sand.	
2046	...do...	21 42 45	158 06 02	119	gy. s.do....	
2047	...do...	21 45 45	158 06 30	217	69	76	gy. and r. s.	...do....	
2048	...do...	21 52 45	158 07 20	1,483	70	76	C.do....	A small man- ganese con- cretion.
2049	...do...	21 59 30	158 08 30	2,226	68	75	br. m.	Red clay...	
2050	...do...	22 07 00	158 10 00	2,555	70	75	br. m.do....	
2051	Jan. 28	22 13 00	158 26 30	2,616	70	75	br. m.do....	
2052	...do...	21 59 30	158 39 00	1,184	70	75	gy. s.	Globigerina ooze.	
2053	...do...	21 59 20	158 25 40	1,651	75	75	gy. s.do....	
2054	...do...	21 48 40	158 17 45	1,237	72	75	gy. m. bk. G	...do....	No specimen.
2055	...do...	21 45 30	158 25 45	536	75	76	gy. m. bk. s.	Globigerina ooze.	
2056	...do...	21 45 20	158 39 00	541	75	76	br. s. G	...do....	Large man- ganese con- cretion.
2057	...do...	21 38 40	158 31 20	440	75	77	br. and bk. s.	...do....	No specimen.
2058	...do...	21 36 40	158 43 00	1,058	75	77	gy. s.do....	Do.
2059	...do...	21 46 15	158 49 20	677	69	76	gy. and bk. s.	Globigerina ooze.	
2060	...do...	21 21 30	158 35 00	1,416	70	75	gy. m.do....	
2061	...do...	21 10 00	158 21 20	1,670	69	75	gy. m.do....	
2062	...do...	21 15 45	158 14 30	952	70	74do....	Do.
2063	...do...	21 03 30	158 01 30	437	69	74	Globigerina ooze.	Small speci- men.
2064	...do...	21 00 20	158 02 00	1,355	69	74do....	
2065	Jan. 29	21 06 40	158 01 00	294	69	74	gy. m.do....	
2066	...do...	21 08 45	157 58 00	278	69	74	gy. m.do....	
2067	...do...	21 10 30	157 57 00	323	69	75	gy. m. and s.	...do....	Manganese nodules.
2068	...do...	21 11 30	157 56 00	307	69	75	gy. m. and s.	...do....	
2069	...do...	21 12 45	157 55 00	287	gy. m. and s.	...do....	
2070	...do...	21 13 45	157 54 30	285	gy. m. and s.	...do....	Pteropods.
2071	...do...	21 14 40	157 53 40	271	gy. m. and s.	...do....	
2072	...do...	21 15 40	157 53 45	201	C.	Coral sand.	Fragments of coral.
2073	...do...	21 16 20	157 54 00	33	C.do....	
2074	...do...	21 16 40	157 53 20	22	C.do....	Honolulu.

1. The first part of the document is a list of names and dates.

2. The second part of the document is a list of names and dates.

3. The third part of the document is a list of names and dates.

4. The fourth part of the document is a list of names and dates.

5. The fifth part of the document is a list of names and dates.

6. The sixth part of the document is a list of names and dates.

7. The seventh part of the document is a list of names and dates.

8. The eighth part of the document is a list of names and dates.

EXPLANATION OF PLATES.

PLATE I.

- Fig. 1. Station 385. 720 fathoms. Coarse Globigerina Ooze.
Orbulina universa d'Orbigny, *Globigerina conglobata* Brady, *Sphaeroidina bulloides* d'Orbigny, *Candeina nitida* d'Orbigny, *Pulvinulina menardii* d'Orbigny, *P. tumida* Brady, *P. micheliana* d'Orbigny.
Magnified 15 diameters.
- Fig. 2. Station 385. 720 fathoms. Fine Globigerina Ooze.
Mostly *Globigerina bulloides* d'Orbigny, with fragments of *Orbulina universa* d'Orbigny.
Magnified 15 diameters.

PLATE II.

- Fig. 1. Station 645. 1,102 fathoms.
Siliceous casts of foraminifera, after treatment with hydrochloric acid.
Magnified 15 diameters.
- Fig. 2. Red Clay Sediment.
Manganese concretions, volcanic sand, crystals of phillipsite, tooth from the lingual ribbon of a mollusk.
Magnified 15 diameters.

PLATE III.

- Fig. 1. Station 688. 1,346 fathoms.
Manganese-iron concretions.
Magnified 15 diameters.
- Fig. 2. Station 338. 2,128 fathoms.
Stellate crystals and spherules of phillipsite.
Magnified 15 diameters.

PLATE IV.

- Fig. 1. Station 670. 1,376 fathoms.
Dark brown, translucent glass, from volcanic mud.
Magnified 15 diameters.
- Fig. 2. Station 995. 2,091 fathoms.
Filamentous, colorless volcanic glass.
Magnified 15 diameters.

PLATE V.

- Fig. 1. Station 746. 2,788 fathoms. Diatom Ooze.
Coscinodiscus rex Wallich.
Magnified 15 diameters.

- Fig. 2. Station 746. 2,788 fathoms. Diatom Ooze.
Segment of valve of *Coscinodiscus rex* Wallich.
Magnified 180 diameters.
- Fig. 3. Station 746. 2,788 fathoms. Diatom Ooze.
Portion of band connecting the valves of *Coscinodiscus rex* Wallich.
Magnified 180 diameters.

PLATE VI.

Diagram of the survey.

PLATE VII.

Track chart, Hawaiian Islands to Midway Islands.

PLATE VIII.

Contour chart, Hawaiian Islands to Midway Islands.

PLATE IX.

Track chart, Midway Islands to Guam.

PLATE X.

Contour chart, Midway Islands to Guam.

PLATE XI.

Track chart, Guam to Luzon.

PLATE XII.

Contour chart, Guam to Luzon.

PLATE XIII.

Track chart, Guam to Yokohama.

PLATE XIV

Contour chart, Guam to Yokohama.



1



2

GLOBIGERINA OOZE.

FOR EXPLANATION OF PLATE SEE PAGE 61.




1



2

1. SILICIOUS CYSTS OF FORAMINIFERA.
2. RED CLAY SEDIMENT.

FOR EXPLANATION OF PLATE SEE PAGE 61.




1




2

1. MANGANESE-IRON CONCRETIONS.
2. CRYSTALS AND SPHERULES OF PHILLIPSITE.

FOR EXPLANATION OF PLATE SEE PAGE 61.



1




2

VOLCANIC GLASS.

FOR EXPLANATION OF PLATE SEE PAGE 61.





AUXILIARY COLLECT

To avoid fine, this book should
or before the date

82945
of U. S. National Museum. No. 55

DATE

NAME

DATE

'41 FAG

82945

ed on

v

10M

